

Pattern recognition in the nucleation kinetics
of molecular self-assembly
Supplementary Information and Data Appendix

Constantine Glen Evans, Jackson O'Brien, Erik Winfree, Arvind Murugan

For arXiv preprint; compiled on July 13, 2022.

Contents

1 Nucleation models and pattern recognition training	7
1.1 The kinetic Tile Assembly Model and Xgrow	8
1.1.1 Estimates of kTAM parameters for SSTs and double-crossover tiles	9
1.2 Stochastic Greedy Model of nucleation	11
1.2.1 Trajectories	13
1.2.2 Nucleation rate	14
1.3 Simplified modes of depletion and winner-take-all effects	17
1.4 Pattern recognition training	22
1.5 Window Nucleation Model	23
1.6 Training and testing image sources and processing	24
1.7 Mixes for pattern recognition experiments	26
2 Fluorophore labels	29
2.1 Motivation	29
2.2 Initial design	29
2.3 Final design	31
3 Tile sequences and layout	35
3.1 Sequences	35
3.2 Fluorophore- and quencher-modified strands	51
3.3 Shape layouts	52
4 Flag patterns	55
4.1 Protocols	55
4.1.1 Flag patterns, constant temperature	55
4.1.2 Flag patterns, ramp	56
4.2 Nucleation model summaries	57
4.3 Individual results	58
4.3.1 H flag 1	58
4.3.2 H flag 2	59
4.3.3 H flag 3	61
4.3.4 H flag 4	62
4.3.5 H flag 5	63
4.3.6 H flag 6	64
4.3.7 H flag 7	66
4.3.8 H flag 8	67
4.3.9 H flag 9	68
4.3.10 H flag 10	69
4.3.11 H flag 11	71
4.3.12 H flag 12	72
4.3.13 A flag 1	73
4.3.14 A flag 2	75
4.3.15 A flag 3	76
4.3.16 A flag 4	78
4.3.17 A flag 5	79

4.3.18 A flag 6	80
4.3.19 A flag 7	81
4.3.20 A flag 8	82
4.3.21 A flag 9	83
4.3.22 A flag 10	85
4.3.23 A flag 11	86
4.3.24 A flag 12	87
4.3.25 M flag 1	89
4.3.26 M flag 2	90
4.3.27 M flag 3	91
4.3.28 M flag 4	92
4.3.29 M flag 5	93
4.3.30 M flag 6	95
4.3.31 M flag 7	96
4.3.32 M flag 8	97
4.3.33 M flag 9	98
4.3.34 M flag 10	100
4.3.35 M flag 11	101
4.3.36 M flag 12	102
4.3.37 M flag 13	103
4.3.38 Uniform concentrations (50 nM)	105
4.4 Plate-level fluorescence data used in Section 4.3	106
5 Pattern recognition	111
5.1 Protocol	111
5.2 Nucleation model summaries	112
5.3 AFM shape counts summary	112
5.3.1 Counts averaged across co-authors and images, normalized to shapes per 25 μm^2 :	113
5.3.2 Counts averaged across co-authors, images and samples, normalized to shapes per 25 μm^2 :	114
5.4 Individual fluorescence and AFM results	115
5.4.1 Hodgkin	115
5.4.2 Avogadro	116
5.4.3 Mitscherlich	117
5.4.4 Hopfield	118
5.4.5 Abbott	119
5.4.6 Moser	120
5.4.7 Horse	121
5.4.8 Anchovy	123
5.4.9 Mockingbird	124
5.4.10 Hazelnuts	126
5.4.11 Apples	127
5.4.12 Magnolia	128
5.4.13 Harom	130
5.4.14 Aon	131
5.4.15 Mbili	133
5.4.16 LetterH	135
5.4.17 LetterA	137
5.4.18 LetterM	138
5.4.19 NoisyApple1	139
5.4.20 NoisyApple2	140
5.4.21 NoisyApple3	141
5.4.22 NoisyHorse1	142
5.4.23 NoisyHorse2	143
5.4.24 NoisyHorse3	144
5.4.25 ObscMagn1	145
5.4.26 ObscMagn2	146

5.4.27 ObscMagn3	147
5.4.28 ObscHazeln1	148
5.4.29 ObscHazeln2	149
5.4.30 ObscHazeln3	150
5.4.31 VarHarom1	151
5.4.32 VarHarom2	152
5.4.33 VarHarom3	153
5.4.34 VarHarom4	154
5.4.35 VarHarom5	155
5.4.36 VarHarom6	156
5.4.37 Uniform concentrations (60 nM)	157

Section 1

Nucleation models and pattern recognition training

Models of nucleation and growth in self-assembling systems must strike a balance between a closeness to physical reality that allows them to capture details of complex behaviors, and an abstraction that allows broader theoretical reasoning. To investigate different questions about our system, we used four different models of nucleation at different points in this spectrum.

More detailed and physically realistic than any of the models we used would be atomic-level¹ and coarse-grained² molecular dynamics simulations of the DNA molecules themselves. Although we do not consider any such models explicitly here, we make use of conclusions derived from such studies by others.³

The lowest-level model we use directly is the single-assembly kinetic Tile Assembly Model (kTAM),⁴ for which a stochastic simulator, Xgrow, is available (Section 1.1). This model abstracts the details of DNA tiles – whose implementations involve strands that interact through hybridization and folding processes – into abstract tiles that attach and detach in single steps and form perfectly rigid lattices. We use the kTAM to analyze the growth of assemblies in our system, and the potential for the formation of chimeric assemblies. To bring the simulation predictions into the ballpark of experimental reality, we attempted to tune the model parameters to fit the single-stranded tile (SST) motif, using experimental and computational results on DNA tile energetics.^{3,5,6}

While simulations in the single-assembly kTAM are useful for favorable growth processes, where assemblies are likely to grow through similar pathways into similar structures and thus the simulation of one, or a few, individual assemblies can be instructive, nucleation involves the exploration of many unfavorable pathways by many small assemblies at once, many of which will melt, and so direct simulation of nucleation by the single-assembly kTAM would be computationally expensive. However, the model was generalized by Schulman & Winfree to consider multiple assemblies growing according to mass-action kinetics; they used this to prove theoretical bounds on nucleation rates that can be calculated with some knowledge of the energy landscape of a system’s possible assemblies and assembly pathways.⁷ To sample assembly pathways for this bound without needing to exhaustively enumerate them, which would be computationally intractable, we developed the Stochastic Greedy Model, which seeks to find likely assembly pathways for assemblies in our system through greedily making favorable attachments and stochastically choosing unfavorable ones. The assembly pathways thus enumerated are then evaluated with respect to the reactions in the generalized kTAM, allowing estimation of pattern-dependent differences in nucleation rates. We used the Stochastic Greedy Model (Section 1.2) for predicting pattern-dependent nucleation rates in our experiments (Figures 3, 4, and 5) as well as for final optimization of pixel-to-tile mapping for pattern recognition.

However, as a stochastic model requiring the collection of many trajectories to produce low-variance estimates of nucleation rates, the Stochastic Greedy Model was either too slow (with more trajectories) or too noisy (with fewer trajectories) to provide the fast, stable comparisons of nucleation rates between similar concentration patterns needed for optimization algorithms searching within the large space of pixel to tile mappings. For this purpose, we instead developed a simple, deterministic, heuristic model, the Window Nucleation Model (Section 1.5), based on intuitive reasoning about the thermodynamics of small assemblies and potential critical nuclei along assembly pathways. This model does not provide a nucleation rate, but instead provides a score value for each concentration pattern such that, if one pattern has a higher score than another, then that first pattern is likely, though not guaranteed, to have a higher nucleation rate in the Stochastic Greedy Model than the second. Using only simple functions on arrays, the calculations for this model were used for the optimization of pattern-to-tile mappings in our pattern recognition experiments, starting from random assignments; once optimization with the Window


Nucleation Model was ended, further optimization was performed with the Stochastic Greedy Model, to fine-tune the pixel-to-tile map and to verify nucleation rates under that more physically-meaningful model.

The Stochastic Greedy Model and Window Nucleation Model, as well as kTAM simulations in Xgrow, consider monomer tile concentrations as being constant over time and not being depleted by assembly, as though controlled by a chemostat. In reality, monomer concentrations deplete as structures nucleate and grow, potentially changing nucleation and growth rates and leading to a winner-take-all effect seen in experiments. We build a simplified model of such a winner-take-all phenomena using a simplified 1-dimensional Markov chain model with representing nucleation and growth, modeled with time-dependent transition rates (Section 1.3).

1.1 The kinetic Tile Assembly Model and Xgrow

The kinetic Tile Assembly Model (kTAM) abstracts DNA tiles as oriented squares with labeled sides.^{4,8} It assumes topologically correct self-assembly, so an assembled object is a connected arrangement of tiles on the square lattice. Tiles are “connected” if they have abutting sides, in which case we may speak of the “bond” between those tiles, whose strength may depend on the pair of labels indicating bond type.

Thermodynamics is specified by associating an energy to each possible assembly based on the tiles involved and how well their side labels match; kinetics is specified by associating rates for each possible tile addition or removal, satisfying detailed balance. Association between two multi-tile assemblies will not be considered in the basic kTAM model, nor will splitting of an assembly into two multi-tile assemblies. (Such interactions, along with lattice defects, can be important for DNA tile assembly,^{9,10} but we did not encounter strong evidence for them in this work.) Thus, considered as a chemical reaction network (CRN) with infinitely many possible species, the reactions are of the form:



where A and A' are assemblies, t_i is a tile of type i , and A' is the same as A but for the addition of tile t at some location. The base model only considers monomer addition reactions, which must be reversible in order to satisfy physical detailed balance, and thus single tile detachments that would break the assembly are not considered. Except when explicitly specified otherwise, kTAM models do not incorporate any notion of globally correct tile placement; they allow any tile to bind in any location so long as at least one bond matches. Thus, tile attachments and assemblies that contain errors with respect to the target complete structure will be considered.

The simplest model for the thermodynamic energy of an assembly considers just the summed interaction strength of all side-to-side bonds based on whether the labels match, and assumes identical bonding strength for matching labels with non-matching labels contributing nothing:

$$G_{bonds}(A) = B G_{se}$$

where B is the total number of matching bonds and G_{se} is the generic bond strength for a suitable choice of units. With respect to these units, detailed balance is expressed, for equilibrium concentrations, as

$$\frac{[A][t_i]}{[A']} = \frac{k_r}{k_f} = \hat{u}_0 e^{-b G_{se}}$$

where b is the number of new matching bonds in A' that were not already in A , and \hat{u}_0 is the reference concentration for these units. We will see later that \hat{u}_0 must be chosen carefully, and will most likely not be standard units. Note that $G_{se} \geq 0$ and more positive values correspond to stronger bonds.

Our simplifying assumption for kinetics is that k_f does not depend on assembly size or tile type. Thus it can be considered as a parameter, and k_r can be calculated for a given tile dissociation from a given assembly based on the detailed balance equation.

Further, it is often simpler and sufficiently insightful to consider assemblies growing in a bath of tiles whose concentrations are held constant. (This scenario is relevant, for example, during the time before significant nucleation has depleted the monomer concentrations.) Here, we will need to allow different tile types to have different concentrations, and we use the notation

$$[t_i] \stackrel{def}{=} c_i = \hat{u}_0 e^{-G_{mc}^i} \tag{1.1}$$

where G_{mc}^i is the parameter controlling the concentration of tile type t_i , with larger values corresponding to lower concentrations (and thus greater positional entropy per molecule). With tile monomer concentrations held constant,

telescoping detailed balance immediately gives us the equilibrium concentration for each assembly:

$$[A]_{eq} = \hat{u}_0 e^{G_{bonds}(A)} \prod_{i \in A} \frac{c_i}{\hat{u}_0} = \hat{u}_0 e^{B G_{se} - \sum_{i \in A} G_{mc}^i} \stackrel{def}{=} \hat{u}_0 e^{-\hat{G}(A)} \quad (1.2)$$

where the assembly energy

$$\hat{G}(A) = \sum_{i \in A} G_{mc}^i - B G_{se} \quad (1.3)$$

is a chemical potential with respect to the tile monomer concentrations.

Because our model does not incorporate interactions between assemblies, each assembly's behavior is independent from the others in the powered scenario, so a single assembly's growth can be modeled as a continuous-time Markov chain (CTMC) where each state is a possible assembly, transitions correspond to the addition or removal of a single tile t_i that is connected by b bonds, and transition rates are given by:

$$r_f = k_f c_i = k_f \hat{u}_0 e^{-G_{mc}^i} \quad (1.4)$$

$$r_{r,b} = k_r = k_f \hat{u}_0 e^{-b G_{se}} \quad (1.5)$$

With every reaction being reversible, this CTMC also satisfies detailed balance, and if it reaches equilibrium (which is only possible for parameters below the melting temperature) then the probability of observing assembly A is:

$$p_{eq}(A) = \frac{1}{Z} e^{-\hat{G}(A)} \quad \text{where } Z = \sum_A e^{-\hat{G}(A)}. \quad (1.6)$$

Note that if a tile is removed from an assembly of size 2, then either tile can be considered the 'remaining' single-tile assembly. As mentioned above, tile removals that would result in multiple, disconnected assemblies cannot be allowed if detailed balance is to be respected.

This simple model can be efficiently simulated by the stochastic Gillespie algorithm^{11,12} using a k -d tree to store rates for log-time steps, as done in Xgrow.¹³ The Xgrow simulator has options to allow a limited variety of reverse reactions that break the assembly into multiple pieces, at the cost of also breaking detailed balance. Specifically, the 'chunk fission' option can be roughly described as: (a) if after a single tile is removed, the remaining assembly is not connected, an irreversible reaction that removes this tile and keeps only the largest remaining assembly is used, with a rate $k_{r,b}$ set by the bonds broken when removing just that tile, and (b) multi-tile subassemblies up to size 2×2 are considered for irreversible removal with rate $k_{r,b}$ where now b is the number of bonds along the perimeter of the subassembly. Without this option, unrealistic surface roughening occurs near the melting temperature due to the inability of small clusters to fall off when they are well-connected internally but have only 1 bond with the main assembly. Although Xgrow does not implement arbitrary fragmentation of large assemblies, during growth of chimeric multifarious structures wherein partial assemblies of two distinct shapes are connected in just one place, circumstances can arise where removal of a single subassembly of size 2×2 or smaller will result separation of the two sides. This is observed in Extended Data Figure E2.

Xgrow has been used extensively in the literature^{9,14,15} as it provides a simplifying clarity, but quantitative comparison with experimental studies of DNA tile systems requires further adjustments, as will be discussed below.⁸

1.1.1 Estimates of kTAM parameters for SSTs and double-crossover tiles

Suppose we have experimentally measured free energies and rate constants for assembly, given for natural units of concentration, time, and energy. We will show that if a few plausible conditions hold, exactly or approximately, then we can find corresponding parameters for the simple kTAM model – thus allowing us use parameters extracted from the literature as a baseline for our simulations of nucleation kinetics.

Specifically, let us assume (1) that the rate constant for tile attachments is identical for all locations and all tile types, so that always $r_f = k_f c_i$ as before; and (2) that the standard free energy of attachment depends only on the the number of matching bonds formed, b , and is linear. Therefore, we can extract the offset and slope, and write:

$$\Delta G_b^\circ = (\alpha - b G_{se}) R T$$

where α and G_{se} are unitless, R is the gas constant, T is the temperature in Kelvin, and the energies use standard units with respect to a standard concentration such as $u_0 = 1$ M.

Considering the per-second rates of tile attachment and detachment at a specific site in an assembly, the simple basic model and the experimentally parameterized model should agree exactly, so

$$r_f = k_f c_i = k_f \hat{u}_0 e^{-G_{mc}^i} \quad (1.7)$$

$$r_{r,b} = k_f u_0 e^{\Delta G_b^\circ / RT} = k_f \hat{u}_0 e^{-b G_{se}} = k_f u_0 e^{-b G_{se} + \alpha} \quad (1.8)$$

where $\alpha = \ln \frac{\hat{u}_0}{u_0}$ and thus the choice of units for the simple model are determined by the offset α via $\hat{u}_0 = u_0 e^\alpha$.

Unfortunately (though not surprisingly) experimental and computational studies^{3,5,6} report that the assumptions we used are only approximately correct for DNA tiles:

1. For DNA double-crossover tiles assembling into lattices on mica, as observed by atomic force microscopy movies,⁵ ΔG_b° was linear in b (within error bars) over a 100-fold range of off-rates, while on-rates were constant within a factor of 2 (except where fewer than 10 data points were collected, e.g., for $b = 4$).
2. For DNA double-crossover tiles assembling in solution, as measured by fluorescence,⁶ a roughly linear dependence on b was also found for ΔG_b° , now covering a 10^9 -fold range of rates, with on-rates remaining constant within a factor of about 4. Importantly, this work also introduced additional loop energy terms that the basic kTAM cannot accommodate, but fortunately these only come into play for “non-convex” assemblies that are rare during nucleation and growth near the melting temperature. In particular, tiles attaching or detaching by 4 bonds involve rare non-convex assemblies, and we are less concerned about inaccuracies of energies and rates for such reactions.
3. For DNA single-stranded tiles assembling in solution, examined computationally by coarse-grained molecular dynamics of 32 distinct local contexts,³ and averaging over contexts with the same b , again we see a roughly linear dependence on b for ΔG_b° covering again a roughly 10^9 -fold range of rates. Although they don’t calculate rates, they too assume a constant for the on-rate. Again, their value for $b = 4$ is anomalous (by $\sim 2 R T$), and again we are content to ignore it.

To find reasonable parameters for the kTAM to model our SST system, we will make use of the computational work³ because the underlying oxDNA model¹⁶ has been well calibrated, and detailed experimental measurements for SST are not yet available. From pages 7-18 of the supplementary materials, the rightmost value of F/kT gives the value of $\Delta G_b^\circ / RT - \ln([t_i]/u_0) \approx 9 - 6 b$ for $b \in \{1, 2, 3\}$. Since $\Delta G_b^\circ = (\alpha - b G_{se})RT$ and in the computational work $[t_i] = 100$ nM and $u_0 = 1$ M, we take $G_{se} = 6$ and $\alpha = -7.1$, and use $k_f = 10^6 / M/s$. Given α and $[t_i]$, G_{mc}^i is determined, completing the parameterization needed for simulation. (Note that recent work¹⁷ with SSTs did not model an α -like effect, resulting in an estimated $G_{se} \approx 8$ for 100 nM tiles that may be too high.)

Note, however, that these parameters are for $T = 50$ °C, as that was the temperature used in the simulations. For our purposes, we are interested in nucleation and growth rates at different temperatures, in particular the critical range between 53 °C and 45 °C. Therefore, we must consider the temperature-dependence of parameters k_f , α , G_{se} , and G_{mc}^i . A roughly 50% increase in the association rate constant was observed⁶ for DNA double-crossover molecules between 12 °C and 21 °C, but we will be satisfied holding k_f constant. We also make the simplifying assumption that α is temperature independent, i.e. that it represents purely entropic factors. Consequently, G_{mc}^i will also be temperature independent.

To provide a temperature dependence for G_{se} , we note that association $b = 1$ corresponds to uncomplicated hybridization of two single-stranded species, for which the thermodynamics is well-understood.¹⁸ In particular, we have

$$\Delta G_b^\circ = (\alpha - G_{se})RT = \Delta H^\circ - T\Delta S^\circ \quad \text{for } b = 1 \quad (1.9)$$

$$G_{se} = \Delta S^\circ / R - \Delta H^\circ / RT + \alpha \quad (1.10)$$

$$T = \Delta H^\circ / (\Delta S^\circ - R(G_{se} - \alpha)) \quad (1.11)$$

We used an implementation of nearest-neighbor energy calculations with SantaLucia 2004 parameters^{18,19} to calculate ΔH° and ΔS° averaged over all binding domain sequences in the SHAM system (excluding the poly-T domains used on shape edges): $\Delta H^\circ = -72.76$ kcal/mol and $\Delta S^\circ = -0.2013$ kcal/(mol· K). For $\alpha = -7.1$, this is effectively linear over the range of interest, with

$$T(G_{se}) \approx 49.74 - 2.85(G_{se} - 5) \text{ °C} . \quad (1.12)$$

For reference, $T = 50$ °C corresponds to $G_{\text{se}} = 4.91$, $T = 48$ °C corresponds to $G_{\text{se}} = 5.61$, and $T = 46$ °C corresponds to $G_{\text{se}} = 6.33$. Concentrations of 16.6 nM, 50 nM and 60 nM correspond to G_{mc} values of 10.81, 9.71 and 9.53, respectively, for pattern recognition, flag, and control experiments.

The above discussion makes it clear that the parameterization of the nucleation model is fairly rough, and we do not expect quantitative agreement with experimental results. Our aim for the model is to provide quantitative insight into the nucleation phenomena with this highly heterogenous system, to observe qualitative effects that may also be present in the real system, and to provide a baseline against which to assess the effectiveness of our sequence design and pixel-to-tile mapping.

With respect to these standard units, we define the free energy (chemical potential)

$$G(A) = \hat{G}(A) - \alpha = -B G_{\text{se}} + \sum_{i \in A} G_{\text{mc}}^i - \alpha \quad (1.13)$$

where as before B is the total number of matching bonds between tiles in the assembly. Now, at equilibrium, with monomer concentrations held constant at their initial values,

$$[A]_{eq} = u_0 e^{-G(A)} = u_0 e^{B G_{\text{se}} - \sum_{i \in A} G_{\text{mc}}^i + \alpha}. \quad (1.14)$$

In the following, we will sometimes find it convenient to refer to a *base concentration* $c = \hat{u}_0 e^{-G_{\text{mc}}} = u_0 e^{-G_{\text{mc}} + \alpha}$ such that individual tiles have concentrations $[t_i] = c_i = c f_i = \hat{u}_0 e^{-G_{\text{mc}}^i} = u_0 e^{-G_{\text{mc}}^i + \alpha}$.

1.2 Stochastic Greedy Model of nucleation

The equilibrium concentration of a particular assembly given by Equation 1.14 does not generally hold while the system is out of equilibrium. In particular, as in the experiments considered here, in an isolated system that initially only contains monomers and does not replenish them as they are incorporated into structures, the total tile concentrations will remain constant, and thus monomer concentrations will deplete. When there is a significant barrier to nucleation, the subset of pre-nucleation structures may quickly approach equilibrium with respect to each other, contributing to some tile depletion, while further growth of nucleated structures continue the depletion.

Nonetheless, Schulman & Winfree [7] show that the equilibrium concentration for an assembly, with respect to the intial monomer tile concentrations, is an upper bound for the concentration of that assembly at any time in the time evolution of a mass-action model with monomer concentrations that deplete during growth, if the initial state consists of monomers and assemblies grow by reversible monomer addition only. As a result, the total rate at which larger assemblies form from a particular assembly A , which we will call its growth flux, has an upper bound of

$$\eta_{\{A\}}^+ = k_f [A]_{eq} \sum_{j \in J(A)} c_j = k_f u_0 e^{-G(A)} \sum_{j \in J(A)} c_j \quad (1.15)$$

where $J(A)$ is the set of possible tile attachments to A and we use that for all times t , $[A](t) < [A]_{eq} = u_0 e^{-G(A)}$.

From this per-assembly growth flux, we can derive an upper bound for nucleation, or more generally, a limiting rate for growth from monomers to a set of particular target assemblies. Consider any set of assemblies S in the space of all possible assemblies, such that every series of tile attachment and detachment events going from an individual tile monomer to the set of target assemblies must contain at least one of the assemblies in S as an intermediate assembly. We might want to consider S to be a ‘boundary surface’ that separates pre-nucleated structures from post-nucleated structures. For example, for a target assembly of size N , the set of all assemblies with size exactly K (for some $K < N$) would satisfy this property. However, in general S can be any set of assemblies that must be crossed during assembly, and this gives us considerable flexibility when finding suitable S . For any such S , considering all series of attachments and detachments, the total flux through the ‘surface’ S , and thus from monomers to target assemblies, has an upper bound of

$$\eta^+ = \eta_S^+ = \sum_{A \in S} \eta_{\{A\}}^+ \quad (1.16)$$

This result is considered more rigorously by Schulman & Winfree [7]. It relies only on S containing a complete surface that all growth must pass through. A poor choice of surface (or a set with extraneous members) will simply result in an unnecessarily high upper bound: for example, in conditions where $G(A)$ does not begin to decrease

until assembly sizes much larger than dimers, the set of all dimers D would still result in η_D^+ constraining the total rate of target production, but the bound would be very high. Generally, a surface that all growth must pass through where $G(A)$ is high for each $A \in S$ will tend to result in the smallest η_S^+ , and thus the tightest bound on nucleation. It would be desirable to find an optimal surface, i.e. one that strictly minimizes η_S^+ .

Even in cases where there are few tile types, regular periodic order, and homogeneous concentrations, enumeration of a complete minimal surface may be infeasible, but excellent approximations can be articulated by taking advantage of symmetry and considering just the most dominant terms. For example, in such cases S may be all assemblies of a well-chosen critical size K , for which the dominant species may be easy to describe. Specifically, as assemblies with higher $G(A)$ within a set S will contribute exponentially less to η_S^+ , it often suffices to consider the assemblies within the set with the *smallest* $G(A)$ values, which will contribute the most to η_S^+ . For homogeneous systems, these will simply be those that have the most bonds with the fewest tiles, often corresponding geometrically to having the largest area for the smallest perimeter.

In the case of unequal concentrations and inhomogenous designs, however, symmetry breaks. Each possible assembly must be considered separately, as each tile may have a different concentration, resulting in a different $G(A)$ despite geometric similarity, and the incorporation of more high concentration tiles at the expense of increasing perimeter may result in $G(A)$ being smaller, breaking the link between geometric compactness and free energy. As a result, simple approaches to finding the largest contributors to a particular η_S^+ , for example, considering squares of increasing size in a homogeneous, single-tile-type system,⁸ cannot be easily adapted to a non-homogeneous system with many tile types.

More generally, consider the set of all possible trajectories from a single tile to a full assembly by monomer addition, and collect – without duplication – the highest-energy point along each trajectory, which we may call the ‘critical nucleus’ for that trajectory. (Using ‘critical nucleus’ for these assemblies is a bit of an abuse of terminology, but please bear with us.) The set S of all such critical nuclei contains a complete separating surface (along with much else) and its lowest-energy point must be reached or exceeded by any trajectory. Pruning this conceptual S down to a smaller subset that contains just enough of the lower-energy assemblies could provide a tractable approximation for the nucleation rate.

Thus, we developed the *Stochastic Greedy Model* to stochastically generate a set S of ‘critical nuclei’ for the nucleation of a single shape, essentially considering the tile system as a uniquely-addressed tile system assembling that one shape. Only correct tile attachments on the uniquely-addressed lattice are considered: attachments of incorrect tiles by one bond are ignored, as are any attachments of tiles that aren’t present in the shape, and each tile i will only appear at a particular site (x, y) on the lattice. The algorithm functions by taking some series of probabilistic trajectories using a ‘two step’ process for each change in assembly: taking an unfavorable step probabilistically, and then deterministically taking the resulting series of possible favorable attachments, while recording the free energy of the assembly immediately after the unfavorable step to try to find a critical nucleus.

Given a set S thus enumerated by the Stochastic Greedy Model, the nucleation rate estimated by Equation 1.16 may be above or below the actual nucleation rate for monomer addition assembly in the bulk mass-action kTAM. To the extent that there are gaps in S such that some kTAM assembly trajectories do not go through S , the estimate will be too low. Both having too few sampled trajectories, and considering only greedy trajectories, contribute to this effect. To the extent that some kTAM assembly trajectories go through multiple states in S , the estimate will be too high. Both enumerating a “thick” surface due to stochastic variability, and the fact that kTAM trajectories may reversibly cross a perfect surface multiple times, contribute to this effect. Finally, to the extent that assembly concentrations for the non-equilibrium kinetics remain lower than the equilibrium concentrations given fixed monomer concentrations, the estimate will be too high. Both tile depletion during growth, and the inclusion of “critical nuclei” whose concentrations are drained by nearly-irreversible post-critical growth, contribute to this effect. Accepting these limitations, we characterize the estimated effective nucleation rate, given the enumerated set of critical nuclei S , in terms of an effective ensemble barrier energy G_{ce} such that

$$\eta_S^+ = k_f u_0^2 e^{-G_{ce}} \quad \text{where} \quad G_{ce} = -\ln \sum_{A \in S} \sum_{j \in J(A)} \frac{c_j}{u_0} e^{-G(A)}. \quad (1.17)$$

We can interpret G_{ce} as the macrostate free energy for a set of microstates that consist of the identified critical nuclei together with an arriving tile that has lost its translational entropy by co-localizing at a prospective site of attachment, but which has not yet formed bonds.

Since our multifarious self-assembly design has the property that a pair of tiles that are adjacent (in a given orientation) in one shape are guaranteed to be non-adjacent (in that orientation) in the other shapes, any multi-tile critical nucleus can be expected to grow specifically into the complete shape that it is consistent with. Therefore,

when S was enumerated based on a specific shape, the nucleation rate estimate for that shape can be written

$$\eta_{shape}^+ = \eta_S^+ .$$

While our algorithm is sufficient for the concentration patterns used in this work, more subtle concentration patterns that are nearly equally localized in multiple structures will require solving the issues raised here. One approach to addressing many of these issues is to adapt a transition path sampling method,²⁰ like forward flux sampling²¹ as a wrapper around the kinetic Tile Assembly Model described above. These statistical sampling methods are explicitly designed to determine the rate of processes controlled by rare events and have been applied to nucleation,²² albeit not in the multi-component limit with unequal concentrations needed here.

We now provide details of the Stochastic Greedy Model.

1.2.1 Trajectories

For each target shape (H, A, M), we generated 40,000 trajectories of possible assembly paths from a single tile all the way to a complete shape. Each trajectory starts from a single tile, used as the initial seed for growth, that is chosen probabilistically by concentration, such that the probability of choosing tile i is $P_{start}^i = c_i / (\sum_i c_i)$. The trajectory will exclusively involve tile additions, such that assembly A_0 is just tile t_i and after s tile addition steps, assembly A_s consists of $(s + 1)$ tiles in total and has free energy (chemical potential) $G(A_s)$ as defined above.

At each point in trajectory construction, where A_s is the current assembly, the algorithm calculates $\Delta G_{att}(x, y) = G(A'_s) - G(A_s)$ for every point (x, y) , where A'_s reflects the potential addition of the correct tile at position (x, y) . We further calculate a Boltzmann weight $p_{att}(x, y) = e^{-\Delta G_{att}(x, y)}$, which we use for the probability that the algorithm will select this site for the next tile attachment. For speed, this calculation is done by keeping values in two arrays, and only changing values adjacent to the change in the assembly array: i.e., for every tile attachment, these two values are recalculated for whichever of the four locations adjacent to that tile attachment are empty. For a given site (x, y) where tile i can attach and form b bonds, we use $\Delta G_{att}(x, y) = G_{mc} - b G_{se} - \ln f_i$ if $b > 0$, while $\Delta G_{att}(x, y) = \infty$ and thus $p_{att}(x, y) = 0$ if $b = 0$.

For the first (presumed unfavorable) step, and subsequently whenever the assembly is again in a state where only $\Delta G_{att} > 0$ attachments are possible, the algorithm chooses a step to take with a probability of attachment $P(x, y) = p_{att}(x, y) / \sum_{X,Y} p_{att}(X, Y)$. Note that this is not the actual probability of that tile attachment taking place in the kTAM, which would be determined only by tile concentrations. If there is no attachment possible at all, i.e. the assembly is complete, then the trajectory ends.

After selecting (x, y) , the algorithm makes the selected attachment (the first of the step's two parts) and updates the ΔG_{att} and p_{att} arrays. The G of the assembly is stored as $G(s)$, where s is the step number. Note that this is stored after the $\Delta G_{att} > 0$ (unfavorable, and probably $b = 1$) attachment. The maximum value of $G(s)$ is stored as the current best guess for the critical nucleus free energy G_{cn} along the trajectory, and the assembly state that resulted in that maximum is stored until replaced by a higher- $G(s)$ assembly.

After the $\Delta G_{att} > 0$ attachment, the algorithm runs a filling routine that repeatedly attempts to make an arbitrary $\Delta G_{att} < 0$ attachment, until no such attachments are possible. Because of the properties of the assembly (all correct tile attachments, only one possible tile per site, no detachment, no negative glues), the order of attachments is not important. This is the second part of the step: making every possible subsequent favorable step after the initial unfavorable step.

As used in this work, the algorithm always continues until the final complete shape is formed and thus no more correct attachments are possible; however, for increased efficiency, it is also defined such that it may stop after a given number of tiles have formed, which the user may believe to be sufficient to guarantee nucleation has occurred.

At the end of a trajectory, the algorithm checks to see whether the final assembly free energy $G(\text{end}) < G(\text{base})$. In other words, it checks whether the final assembly is more favorable than a low-concentration monomer tile, for equilibrium with respect to fixed tile concentrations. If this is not the case, then the algorithm reports the trajectory as a failure. As used in this work, where trajectories are computed all the way to the complete shape, either all trajectories are successful or all trajectories are failures, given a concentration pattern, temperature, and shape. With respect to the fixed tile concentrations, we define the melting temperature for a given pattern and shape as point where $G(\text{complete}) = G(\text{base})$. (Melting temperatures would not be pattern-dependent if all tiles were shared in all shapes, but as it is, different patterns may place more high-concentration tiles in different shapes; the additional shape-dependence comes from the different area-to-perimeter ratios.)

The trajectory algorithm then returns the following information to the nucleation rate algorithm:

- The G_{cn} found by the algorithm: the highest $G(s)$ along the trajectory.

- The critical nucleus assembly that corresponds to that G_{cn} .
- $G(s)$ for every step s .
- The step s where G_{cn} is found.
- Whether $G(\text{end}) < G(\text{base})$.

Example greedy trajectories for H flag 6 are shown in Fig. 3c. Figure 1.1 shows a Boltzmann-weighted sample of critical nuclei in the enumerated sets S for each shape, for the H flag 6 concentrations at $T = 49.2^\circ\text{C}$, at $T = 48.3^\circ\text{C}$, and at $T = 45.5^\circ\text{C}$. Given the quantitative differences between the theoretical model and experimental results, we only look to interpret qualitative features of the critical nucleus sets. It is immediately clear that (a) most critical nuclei in H are near the flag, while critical nuclei in the other shapes are scattered around; and (b) the typical critical nucleus size is strongly dependent on temperature in the samples shown, varying from 3 to 16 for the on-target shape and from 3 to 73 for the off-target shapes, with the greatest difference being in the off-target shapes. Figure 1.2 shows an analogous Boltzmann-weighted sample of critical nuclei for the Avogadro pattern recognition experiment. The phenomena are similar here, although the on-target critical nuclei are somewhat larger on average, off-target critical nuclei are somewhat smaller, and estimated nucleation rates are somewhat closer to each other, consistent with the narrower range of temperatures used. As the critical nuclei for the on-target shape can be quite small, there is a sense in which the selectivity of pattern recognition relies on the rejection of nucleation attempts in the off-target shapes, where the amount of information being considered is governed by the size of their critical nuclei at the relevant temperature.

Fig. 3e shows macrostate ensemble free energy for assemblies of a given size, using the ensemble of trajectories generated by the Stochastic Greedy Model. Let S_K be the set of size- K assemblies that appear in some enumerated trajectory. Then we calculate and plot $G_{SGM}(K) = -\ln \sum_{A \in S_K} e^{-G(A)}$. Because the greedy trajectories are biased toward the low-energy assemblies that dominate this Boltzmann-weighted sum, G_{SGM} is an adequate approximation of the macrostate ensemble free energy for the set of *all* size- K assemblies for the given shape.

1.2.2 Nucleation rate

The Stochastic Greedy Model generates a set number of trajectories. Once these trajectories have been generated, the algorithm has a set of distinct critical nuclei, S , and associated information. Using the enumerated set S , the algorithm estimates the ensemble nucleation barrier energy G_{ce} and the nucleation rate η^+ according to equations 1.15, 1.16, and 1.17, with one adjustment: rather than summing over all possible tile additions from each critical nucleus, $J(A)$, we sum over only the favorable (energetically downhill) tile additions, which we call $J^+(A)$. While the full set $J(A)$ is required for a rigorous upper bound in the general case, for the set of critical nuclei enumerated by the Stochastic Greedy Model, which are guaranteed to have at least one downhill tile addition step available, neglecting the uphill tile additions avoids counting flux that is likely to be immediately reversed, while keeping the dominant favorable tile additions. Thus, we *replace* the previous formulas with:

$$\eta_{\{A\}}^+ = k_f [A]_{eq} \sum_{j \in J^+(A)} c_j = k_f u_0 e^{-G(A)} \sum_{j \in J^+(A)} c_j \quad (1.18)$$

$$\eta^+ = \eta_S^+ = \sum_{A \in S} \eta_{\{A\}}^+ = k_f u_0^2 e^{-G_{\text{ce}}} \quad (1.19)$$

$$G_{\text{ce}} = -\ln \sum_{A \in S} \sum_{j \in J^+(A)} \frac{c_j}{u_0} e^{-G(A)}. \quad (1.20)$$

The total nucleation rate for a shape gives no information on the region where nucleation began, even though, with unequal tile concentrations, some regions may be much more likely than others to participate in nucleation rather than to grow by tile attachments after a structure has nucleated. While trajectories in the Stochastic Greedy Model start from individual tiles, the trajectories are used to find distinct critical nuclei, and a particular critical nucleus may arise from trajectories starting from many different positions. An alternative measure of the contribution of a particular location to the nucleation rate of a shape, calculated only from the set of distinct critical nuclei S that was enumerated for the given shape, is the sum over all critical nuclei that include the given location, of the nucleation flux through the nucleus divided by its size:

$$\eta_{\text{shape}:x,y}^+ = \sum_{A \in S \text{ s.t. } x,y \in A} \eta_A^+ / N(A) \quad (1.21)$$



Figure 1.1: A Boltzmann-weighted sample of 6 ‘critical nuclei’ for each shape, at three temperatures, for H flag 6. Orange and red indicate low- and high-concentration tiles that are part of the critical nucleus, respectively; grey and green are used analogously for tiles not in the nucleus. η^+ is the estimated nucleation rate for the shape; G_{ce} is the corresponding critical nucleus ensemble free energy; G_{se} is per-bond differential sticky-end energy at the shown temperature; and G_{cn} is the chemical potential of the given critical nucleus.




Figure 1.2: A Boltzmann-weighted sample of 6 ‘critical nuclei’ for each shape, at three temperatures, for Avogadro. The color scale from orange to red scales linearly from the lowest to highest concentration, and is used for tiles that are part of the critical nucleus; the color scale from black to white is used analogously for tiles not in the nucleus. η^+ is the estimated nucleation rate for the shape; G_{ce} is the corresponding critical nucleus ensemble free energy; G_{se} is per-bond differential sticky-end energy at the shown temperature; and G_{cn} is the chemical potential of the given critical nucleus.

where $N(A)$ is the number of tiles in A . By weighting the nucleation rate in this way, the contributions of regions to the total nucleation rate of a shape can be calculated by adding the contributions of each location, and the sum of contributions for all locations in a shape equals the total nucleation rate:

$$\sum_{x,y \in shape} \eta_{shape:x,y}^+ = \eta_{shape}^+$$

Because we performed a rough calibration of the kTAM parameters in Section 1.1.1, we can use the Stochastic Greedy Model to estimate the temperature dependence of nucleation rates for any given shape and pattern of tile concentrations. Simultaneously, for each temperature we can calculate the Boltzmann-weighted (i.e. weighted proportional to equilibrium concentration) average critical nucleus size. In Sections 4.3 and 5.4, we perform this analysis for each sample from the flag scan and pattern recognition experiments, respectively. These calculations use the initial concentrations of each tile, and do not model the depletion process. Although both nucleation rate and average critical nucleus size can be calculated above the shape- and pattern-dependent melting temperature (where $G(\text{complete}) = G(\text{base})$), their interpretation becomes more fraught, so we mark the melting temperature with a dashed vertical line and plot only at lower temperatures.

It should be emphasized that our nucleation rate estimates are not intended to be quantitatively accurate: the kTAM model and its calibration only roughly capture the thermodynamics and kinetics of our system; the further approximations used in the Stochastic Greedy Model are also rough; and nucleation kinetics are notoriously sensitive with order-of-magnitude differences between theory and experiment being common and often difficult to explain.^{23,24} (Example deficiencies in our estimates can be seen in the plots for flag scan experiments, which contain abrupt jumps in the nucleation rate estimate that are artifacts of the Stochastic Greedy Model’s enumeration process for ‘critical nuclei’.)

In particular, we do not expect the model to accurately predict the temperature (during a ramp experiment) at which nucleation becomes measurable, nor the absolute or relative nucleation rates at a given temperature. For example, in the pattern recognition experiments, the time to go from about 5% to 10% quenched was at least about an hour, indicating that nucleation was never faster than about 5×10^{-13} M/s, and this mostly occurred between 47 °C and 48 °C. The nucleation model, on the other hand, estimated that this rate would occur between 48.4 °C and 48.6 °C for most patterns, and that at this temperature, the on-target shape would nucleate at least 2 (and often more than 5) orders of magnitude faster than the off-target shape – which fails to explain cases such as Mockingbird where the off-target shape nucleates at nearly the same temperature. This suggests that the nucleation rate estimates may be a few orders of magnitude too high; were k_f decreased, the experimentally-observed nucleation rates would correspond to a lower temperature where on-target and off-target rates are more similar in the model.

Nonetheless, our model provides several important insights (perhaps better phrased as hypotheses) that help us appreciate features of nucleation in multifarious self-assembly systems and general characteristics of their ability to perform pattern recognition. Most notable is the sharp increase in nucleation rate for the on-target shape as the temperature drops below its melting temperature, in contrast to the much more gradual increase for the off-target shapes. This is accompanied by a fast transition (i.e. over a narrow temperature range) from large critical nuclei to small critical nuclei for the on-target shape, whereas the transition occurs much more slowly for the off-target shapes.

Consequently, selectivity for the correct target shape (i.e. the ratio of nucleation rates) increases as one considers increasing temperatures, with the best selectivity near the melting temperature where absolute nucleation rates are low. However, consistent with the shapes having different melting temperatures (M has the lowest melting temperature due it having the highest perimeter-to-area ratio), it is possible that as one gets *too* close to their melting temperatures, the target shapes’ nucleation rate may plummet below the off-target shapes (as occurs for all the M flag scan patterns). That said, the relative stability of on-target nucleation rates sufficiently below the melting temperatures provides another perspective on the previously-mentioned interpretation that pattern recognition selectivity comes primarily from rejection of off-target nucleation attempts.

1.3 Simplified modes of depletion and winner-take-all effects

The above model of nucleation in multi-component systems provides a basis for understanding the dependence of nucleation rate on concentration patterns, and thus the ability of self-assembly to perform pattern recognition, under the assumption that the temperature and monomer tile concentrations are fixed. The model considers each shape independently, and thus could be used to ask whether a single uniquely-addressed shape could recognize

patterns via a nucleation rate that is above a “YES” threshold or below a “NO” threshold, depending on whether the pattern should be recognized. To classify patterns into multiple groups (such as H, A, and M) one could use three distinct uniquely-addressed shapes, sharing no tiles, if the patterns contain enough redundant information that a YES/NO decision for each shape is possible based on a partitioning of pixels into three groups. We expect this would be possible for the training set of 18 images used in this work, for example, using 3 new shapes each containing 300 tiles.

What then is the advantage of a multifarious self-assembly system with shared tiles, if any? Intuitively, we expect two advantages. First, a system with fully-shared tiles would enable all shapes to be responsive to all pixels, and thus redundancy in the pattern information would not be necessary. Our system, with a partially shared tile set, partially addresses this issue. Second, when one shape nucleates first, it will deplete the concentrations of the shared tiles, and this may limit the nucleation of other shapes, resulting in a winner-take-all effect. Such a winner-take-all effect allows experiments with a temperature ramp that does not require precise knowledge of the nucleation temperatures for on- and off-target structures. In contrast, in systems without shared components, experiments would have to be carried out at a specific temperature that allows specificity; such a temperature may be hard to know *a priori* and will not be universal, varying with the specific patterns and structures in question.

Here, we wish to quantify this winner-take-all (WTA) effect and provide some understanding of under what circumstances it will or will not occur, and how it can aid the pattern-recognition process. These questions could presumably be addressed by building on the kTAM or SGM, adding equations to model the depletion of each tile type during the nucleation and growth of self-assembling structures, but doing so would be prohibitively expensive computationally. Further, it is hoped that simplifying to the essential ingredients provides more insight.

Absence of winner-take-all in homogeneous systems. Before examining how monomer depletion can lead to a winner-take-all effect in multifarious systems, it is worth commenting on why this effect would not be expected in simpler systems. Crystal polymorphism has been well-studied since the time of Mitscherlich; the ability to nucleate only the most stable form of several polymorphs is an industrially relevant problem, in domains ranging from making chocolate to drugs. Consequently, numerous annealing protocols have been developed to enhance the yield of, say, the most stable polymorph while minimizing nucleation and growth of the competing phases.

The winner-take-all effect described here makes the selectivity problem easier in the multi-component limit and is absent in the well-studied case of homogeneous crystals. In the multi-component structures studied here, the on-target structure, e.g., H in the models above, can nucleate from a region of high concentration tiles while critical seeds of the off-target structure, say A, predominantly involve low concentration tiles (since the same high concentration tiles are not spatially localized on A). Depletion affects the low concentration tiles much more significantly than the high concentration tiles; hence depletion will more significantly affect the nucleation rate of A (the off-target structure) than the nucleation rate of H (the on-target structure). In contrast, in homogeneous crystal polymorphs, depletion will affect nucleation of both on and off target structures to similar extents because both nucleation rates are determined by the concentration of the same component(s). Hence, we do not expect a winner-take-all effect for crystal polymorphic structures involving only one or a few kinds of components.

Presence of winner-take-all in multifarious systems with shared components but not in multi-component systems with independent components. We will consider the assembly of two structures, say ‘H’ and ‘A’, from a pool of monomers that deplete over time as monomers get incorporated into the structures. The structures may be competing for shared monomers, or may not be competing. Two models will be developed. The first is a highly simplified model that just tries to capture the notion of classical Arrhenius nucleation kinetics combined with uniform growth and maturation using shared components. We call it the “coarse-grained” (CG) CRN model. The second model is still oversimplified, but it adopts more details and an overall framework similar to the SGM, with individual tiles and individual assemblies growing (or shrinking) by reversible monomer addition reactions that satisfy detailed balance. We call it the “SGM-lite” CRN model. They are both illustrated schematically in Figure 1.3.

The coarse-grained model. The CG model has two variants, one considering all tiles being shared between the two shapes, and the other considering all tiles to be distinct. The shared-tiles (S) variant has the following 6





Figure 1.3: Schematic illustration of the CG CRN model (top) and the SGM-lite CRN model (bottom). On the left are the two shapes, called “H” and “A”, with black indicating the set of tiles whose concentrations will be enhanced in this study to favor nucleation of H, and intermediate grey indicating another set of tiles that could be used to favor nucleation of A. On the right we indicate the assemblies that are considered in the models: The CG CRN just uses three assemblies for each shape, e.g. H_{nuc} , H_{mid} , and H_{full} . The SGM-lite CRN considers all assemblies that occur in M pre-computed stochastic greedy trajectories for each shape, starting from a single high-concentration tile and proceeding by monomer addition to the full shape. Outlines of sample assemblies along three color-coded trajectories are shown for each shape.

reactions:



where $N = 500$ is the total number of tiles in each shape, $s = 25$ is the number of high concentration tiles and also the number of tiles in the assemblies H_{nuc} and A_{nuc} , and species c_n^H , c_n^A , and c_g respectively represent the tiles in two nucleation regions and the growth region (black, intermediate grey, and light grey in Figure 1.3). Because our reactions represent coarse-grained pathways involving many individual tile addition steps, in each class the concentrations of all tiles will be the same, and we can model it using a single species concentration. That is, the first pair of reactions represent the nucleation and growth of all s tiles in the nucleation regions; the second pair of reactions represent the growth of the assembly to include all but the tiles that could nucleate the opposing shape; and the third pair of reactions completes the shape. Thus, one copy of each relevant tile is consumed in each coarse-grained reaction step.

The rates for these coarse-grained reactions are calculated from approximations of kTAM kinetics incorporating classical nucleation theory. Specifically, we assume that in the nucleation region there is a critical nucleus whose size depends on temperature (via the sticky-end binding strength G_{se}) and tile concentration (via the energy parameter $G_{mc} = \alpha - \ln c/u_0$), and the nucleation rate is proportional to its equilibrium concentration. (As usual, $\alpha = -7.1$ and $u_0 = 1 \text{ M}$.) That is, we take $k_{nuc} = k_n c_{\text{critical}} = k_n u_0 e^{-G(A_{\text{critical}})}$ where $G(A_{\text{critical}}) = -2k(k-1)G_{se} + k^2 G_{mc} - \alpha$ for a $k \times k$ square critical nucleus, and $k = G_{se}/(2G_{se} - G_{mc})$ is determined by maximizing $G(A_{\text{critical}})$ to find the free energy barrier. We interpolate k to continuous real values, rather than integers, for convenience. We set $k_n = 0.01 / \text{s}$ to roughly match SGM nucleation rates for the range of concentrations considered. The time for growth of the rest of the nucleation region is assumed to be negligible with respect to nucleation itself. (Note that H_{nuc} and A_{nuc} do not represent the critical nuclei themselves, but rather the entire region that is considered prone to nucleation.) A weakness of this simple model is that it does not consider entropic effects due to there being many places where nucleation could initiate and likewise many alternative critical assemblies that could contribute to the overall nucleation rate.


For growth of assemblies, our reference rate considers the expected time for a biased random walk to complete N steps, with forward steps occurring at rate $k_f c$ and reverse steps occurring at rate $k_{r,2}$, where as per the kTAM, $k_{r,b} = k_f u_0 e^{-b G_{se} + \alpha}$ and $k_f = 10^6 / \text{M/s}$. That is, it is the net rate for completing the growth of the entire shape assuming each reversible step involves forming or breaking 2 bonds. For coarse-grained steps that involve fewer net steps, the rate is accordingly rescaled. Note that barriers for initiating a new layer of tiles on a facet are not considered here, although they can be very significant near the melting temperature when $2G_{se} \approx G_{mc}$.

Altogether, this gives:

$$\begin{aligned} k_{nuc}(T, c) &= k_n u_0 e^{-G_{se}^2/(2 G_{se} - G_{mc})} && \text{if } 2 G_{se} > G_{mc} \text{ else } 0 \\ k_{grow}(T, c) &= (k_f c - k_{r,2})/N && \text{if } 2 G_{se} > G_{mc} \text{ else } 0 \end{aligned}$$

where G_{mc} is a function of c and G_{se} is a function of the temperature T as per Equation 1.12. We will consider an anneal from well above the nucleation temperature for either shape (for our range of concentrations) down to a temperature where growth is strongly favored, specifically, from 55 °C to 30 °C over some time t_{ramp} during which G_{se} increases linearly with time.

The no-shared-tiles (NS) variant has just the following 4 reactions, as there is no need to distinguish two populations of growth tiles:



Results from simulations of the CG model are shown on the left in Figure 1.4. For comparison, the flag experiments of Figure 4 cooled from 48 °C to 46 °C over 100 hours, for a rate of 0.02 °C per hour; the slowest simulation in the CG model traverses 25 °C over the same time, i.e., it is 12.5 times faster. We consider this range because the CG model’s approximations appear to overestimate reaction rates, making the CG results “too perfect” for slower ramps. Similarly, the fact that the CG model considers high-concentration tiles within a solid region, rather than in a checkerboard pattern, makes the model more sensitive to concentration enhancements, and thus we consider a maximum 6-fold enhancement instead of the 17.6-fold enhancement used in the flag experiments.

The general trends are as expected: For model variant (S) with shared tiles, greater concentration enhancements and/or slower temperature ramps result in both higher selectivity and greater completion. In contrast, while the model variant (NS) with no shared tiles similarly obtains greater completion for greater concentration enhancements and/or slower temperature ramps, it only obtains high selectivity for very fast ramps (for which completion is poor). This is because, without competition for tiles, both shapes will eventually grow to completion in the NS variant of the model; observing the results before it has had time to grow is the only way to obtain selectivity, which results from significant nucleation from co-localized high-concentration tiles occurring at an earlier time in the ramp than nucleation from low-concentration tiles in the off-target shape.

(Note that if we considered a temperature hold instead of a ramp, then we could work at a temperature that is initially very selective for the target shape, if only we knew what temperature that should be. But due to concentration depletion, nucleation and growth would come to a halt before reaching completion.)

Given the simplification inherent in this model, it is worth considering which phenomena are likely to hold up in more accurate models, differing quantitatively but not qualitatively, and which phenomena might change substantially. A few concerns may be considered. The CG behavior for fast ramps may not transfer to the experimental system because: (1) SST form unstructured aggregates rather than well-defined crystalline assemblies when cooled too fast; (2) classical nucleation theory assumes that the critical nuclei are at equilibrium instantly, whereas our experimental results suggests that nucleation and growth kinetics are not well separated. (Specifically, when a single shape contains multiple labels at different locations, experimental fluorescence trajectories commonly exhibit a temporal order of quenching consistent with the distance between the expected nucleating center and the fluorophore position within the shape, indicating that timescales for creating critical nuclei and for completing the growth process are overlapping.) The CG behavior for slow ramps may not transfer to the experimental system because: (1) near the melting temperature, facet nucleation barriers will limit growth, but these are ignored by the model; (2) nucleation and growth are modeled with irreversible reactions, but as concentrations drop, small assemblies will become unfavorable and melt due to Ostwald ripening. To gauge the severity of these issues, we explore a slightly less oversimplified model.





Figure 1.4: Simulation results examining the winner-take-all effect due to depletion during an anneal. The CG model is shown on the left, and the SGM-lite CRN model is shown on the right. All simulations were done with a base concentration of 50 nM. The top plots show concentrations of complete assemblies and nucleation rates as a function of time for an intermediate temperature ramp rate (50 hours and 500 hours respectively) and an intermediate concentration enhancement (3-fold and 9-fold respectively). In the lower plots, selectivity is computed as the fraction of final complete assemblies that are the target shape, H, and completion is computed as the final concentration of the target shape as a fraction of the limiting base concentration. Nucleation is computed as the rate of production of H_{nuc} or A_{nuc} in the CG model, and as the net rate of production of assemblies containing at least 25 tiles in the SGM-lite model.

The more detailed model. The SGM-lite model also has two variants, one considering tiles being shared between the two shapes identically as in the experimental system, and the other considering all tiles to be distinct. All reactions in this CRN are of the form:



where A is an assembly (or single tile), t_i is a single tile, and A' is assembly A with tile t_i added in the correct position with respect to either the on-target or off-target shape, forming $b \geq 1$ bonds. Due to computational constraints, not all such reactions are considered. Rather, for each shape we generate $M = 8$ trajectories starting with a single high-concentration tile, adding a random tile that attaches by at least 2 bonds if that is possible, else adding a random tile that attaches by just 1 bond, until the full shape has been grown. As we use the actual tile arrangement for the experimental H and A shapes, which have respectively 480 and 496 tiles, this leads to roughly 8000 reactions and a comparable number of species. Because nucleation in the model is confined to start with the high-concentration tiles for a particular pattern, in this case H flag 6, the resulting CRN is only suitable for simulating that concentration pattern – but for any level of concentration enhancement and at any temperature. Thus, by letting G_{se} be a function of time, as in the CG model, we may use a standard simulator for mass-action chemical kinetics [25].

Results from simulations of the SGM-lite model are shown on the right in Figure 1.4. The simulated cooling rate is now ten times slower than for the CG model, 0.025 °C/h compared to the experimental 0.02 °C/h, but over a wider range, again from 55 °C to 30 °C.

Many trends are similar as in the CG model, but there are some noticeable differences. Perhaps most noticeable is that in the shared-tile model variant, the yield of complete on-target shapes never exceeds 60% for our parameters, even under circumstances where the selectivity is near perfect. This is related to another striking difference: that the off-target nucleation rates are delayed from but not suppressed by the preceding on-target nucleation and growth – both peak at about 1 nM/h although the off-target nucleation occurs later, unlike in the CG model where the maximum off-target nucleation rate is more than 10-fold lower than the off-target rate. What we observe

happening is that while the on-target shape nucleates and slowly grows to its full size, the off-target shape nucleates but grows to only on average a little less than 100 tiles in size. Sensitivity to slow growth kinetics can also be seen in the surprising reduction of selectivity in the shared-tile model as the ramp gets slower, for low concentration enhancements. Slowing the annealing further by 10-fold only moderately ameliorates this effect.

A fuller picture of the winner-take-all effect in pattern recognition by nucleation in multifarious self-assembly will require further investigation.

1.4 Pattern recognition training

The goal of pattern recognition training is to optimize the pixel-to-tile map θ so as to enhance nucleation of the target shape for each training image, while suppressing nucleation of the off-target shapes. The tile set itself stays the same, so with the SHAM system, the set of images will be classified into three groups according to the resulting shapes H, A, and M. However, generalization to a greater number of classes/shapes is straightforward, just requiring the design of a multifarious system with more possible shapes.

Several decisions must be made in order to numerically define the objective function for optimization. How should nucleation rates be predicted? By what metric should deviations from the ideal be measured? Should additional criteria be incorporated to address experimental concerns? We will use a score based on maximizing the ratio of predicted nucleation rates using a fast and simple approximation, adjusted to impose preferences for locating nucleation near areas that worked well in flag experiments, and to impose penalties against using high concentrations for fluorophore / quencher strands and their neighbors.

The overall score for optimization will be based on a modification of the Stochastic Greedy Model's critical ensemble nucleation barrier energy estimate G_{ce} , simplified and adjusted (using penalties $w_{x,y}$) to predict low nucleation rates for empirically poor areas in each shape. This modification is described below as the Window Nucleation Model. Thus, given a pixel-to-tile map,

$$\text{Score(image, shape)} = G_{ce}^{\approx}(\text{image}, \text{shape}) \quad (1.22)$$

gives a score that is lower for faster-nucleating cases. Since the nucleation rate is exponential in G_{ce} , the difference in scores between the target shape and its closest competitor is the log of the ratio of their nucleation rates. This gives a score for the image, in which we also add penalty for tiles that should remain at low concentrations.

$$\text{Score(image)} = \text{Score(image, target shape)} - \min \text{Score(image, other shape)} + \sum_i d_i f_i \quad (1.23)$$

where $f_i = c_i/c$ is the concentration of tile i relative to the base concentration c , so $1 \leq f_i \leq 27$. This per-image score is also lower for images predicted to perform better. The penalties d_i are detailed below.

Finally, these scores were combined for all 18 training images to give a total score for that assignment map:

$$\text{Score(overall)} = \max_{\text{images}} \text{Score(image)} \quad (1.24)$$

That is, we try to optimize the pixel-to-tile map by improving the worst-performing image, in hope of obtaining a system whose performance is reasonably consistent across the training images.

Specifically, the overall score was minimized using a naive, parallel “hill-climbing” (in our case, hill-descending) algorithm. Moves consisted of attempted swaps of randomly-chosen pairs tiles, accepted when the score decreases. 72 parallel hill-climbing processes attempted to minimize the score, but every 100,000 steps, all hill-climbing processes switched their current state to the best state any of them had found so far.

After minimization with the Window Nucleation Model, we continued with a slower fine-tuning optimization stage using the Stochastic Greedy Model's estimate for G_{ce} instead of G_{ce}^{\approx} . (Thus, the position-dependent $w_{x,y}$ penalties were no longer applied, although the tile-specific concentration penalties d_i were still applied.) At the time of systems design, this process used an older version of model, in which (i) $G_{se} = 7.0$ and $G_{mc} = 12.75$, (ii) G_{ce} was estimated from just 50 trajectories, each of which took only 14 unfavorable steps rather than completing the entire shape, and (iii) there was no cutoff for the final G of the trajectory. Additionally, each parallel hill-climbing process synchronized with others every step, as each step was quite slow so synchronization adds little delay. During the fine-tuning step, ~ 25 tiles were swapped.

To promote consistent levels and behavior of fluorescence signals, we planned to use fluorophore- and quencher-labeled strands at the base (i.e. lowest) concentration for each experiment; therefore, we wanted to restrict the pixel-to-tile map so as to accommodate this choice. The tiles with non-zero tile-concentration penalty d_i were as

follows: locations with fluorophore labels ($d_i = 1,000$), locations with quenchers ($d_i = 1$), and tiles adjacent to fluorophore-quencher pairs that bound to both the fluorophore and quencher ($d_i = 0.5$). Because the system used 917 tiles and 900 pixel locations, there were 17 assignments that were, for any pattern, always at base concentration. Consequently, the 12 fluorophores, with such a high penalty, always ended up with those assignments, while some quenchers and adjacent tiles were not always at base concentration, but were optimized toward the lower end of concentrations.

In experiments, both fluorophore- and quencher-labeled strands were used at the base concentration, regardless of the concentration implied by the corresponding image pixel (but adjacent tiles were used at the pixel-specified concentration). Accordingly, during training the nucleation rate estimates used base concentrations for fluorophore- and quencher-labeled strands, regardless of the value of the pixel indicated by the current pixel-to-tile map (but again adjacent tiles used pixel-specified concentrations).

1.5 Window Nucleation Model

To optimize the pixel-to-tile map for training pattern recognition by nucleation, we need a fast inner loop, and thus it is infeasible to use the full Stochastic Greedy Model. Finding a pixel-to-tile map that will work well experimentally does not necessarily require an accurate nucleation rate estimate; any function that provides a sufficiently similar rank ordering should be usable. Our choice was to use the simpler and faster Window Nucleation Model, which can be interpreted as calculating G_{ce} as in equations 1.18, but for a boundary set of ‘critical nuclei’ S that consist of every $k \times k$ region (‘window’) in the shape, and with the sum over $J^+(A)$ replaced by a single constant concentration, the base concentration c . In that case, we approximate

$$G_{ce} \approx -\ln \sum_{A \text{ is } k \times k} \frac{c}{u_0} e^{-G(A)} = -\ln \sum_{A \text{ is } k \times k} \frac{c}{u_0} e^{\alpha + B G_{se} - \sum_{i \in A} G_{mc}^i} = \text{const} - \ln \sum_{A \text{ is } k \times k} \prod_{i \in A} f_i \quad (1.25)$$

where $c_i = u_0 e^{-G_{mc}^i + \alpha} = c f_i$ and c, B, G_{se} , and α are constants that can be factored out as an additive offset to G_{se} .

The Window Nucleation Model calculates a score related to this approximation for G_{ce} , but also including position-dependent weights $w_{x,y}$ to encourage pattern recognition to make use of tiles that were demonstrated to work well in the flag scan experiments. Given a pixel-to-tile map θ to be evaluated, tile concentrations are stored in a zero-indexed $L \times L$ array as $f(x, y) = c_i/c$ where i is the tile at position (x, y) within the shape, or $f(x, y) = 0$ if the position is outside the shape. The concentration $c_i = c e^{3 p_n \ln 3}$ is determined from the pattern image being evaluated based on $i = \theta(n)$, with p_n being the grayscale level for image pixel n , and the base concentration is $c = 16.67$ nM. Then the score for window size k is calculated as:

$$\text{Score(image, shape)} = -\ln \sum_{X=0, Y=0}^{L-k, L-k} \exp \sum_{x=X, y=Y}^{X+k-1, Y+k-1} w_{x,y} \ln f(x, y) = -\ln \sum_{X=0, Y=0}^{L-k, L-k} \prod_{x=X, y=Y}^{X+k-1, Y+k-1} [f(x, y)]^{w_{x,y}} \quad (1.26)$$

where $w_{x,y}$ is a weight (by default equal to 1) provided for each location in the shape to roughly incorporate experimental observations about which areas nucleated better than others. Because $f(x, y) \geq 1$, a weight $w_{x,y} > 1$ will result in a more negative score, i.e. a prediction of faster nucleation, whenever the relevant tile is above the base concentration. Note that the contribution to the nucleation score is zero for window regions that extend beyond the boundaries of the shape. Also note that temperature, in the form of its influence on G_{se} , does not come into play in this score, as bond energies can be factored out as a constant independent of position and pixel-to-tile map. Because temperature does influence the size of critical nuclei, our choice of k has implications on the range of temperatures for which the Window Nucleation Model will be reasonably well correlated with actual nucleation. The area K discussed in the main text as the size of a putative critical nucleus here corresponds to $K = k^2$.

For pattern recognition training, the window size was $k = 4$. This was chosen to be slightly smaller than the 5×5 flag size, which we knew could be sufficient for selective nucleation, because in the pattern recognition setup the high-concentration tiles need not be restricted to a checkerboard arrangement within the window; however, we make no claims of optimality here. The position-dependent weight was 2 when considering tiles in a target shape for a particular pattern within the 5×5 contiguous region of the following flags: H flags 2, 6, 8, and 10; A flags 1, 9, 10, and 12, and M flags 1, 5, 9, and 11, and was otherwise 1. When considering tiles in an off-target shape, all weights were 1. The goal of this weighting was to push high concentration regions for nucleation of the target shape into regions shown to work well in the flag-scan experiments (Fig. 4d), in the sense of having strong on-target nucleation and growth, but low off-target nucleation and growth.




Figure 1.5: Nucleation heatmaps for the indicated training patterns are superimposed on their respective target shapes. The color of an (x, y) location is the RGB sum of adjusted heatmap greylevels (linearly proportional to the spatial nucleation rate $\eta_{x,y}^+$ for the given shape and pattern, but adjusted for visual clarity) times the given pattern’s indicated color. Colored outlines indicate the smallest region accounting for 75% of nucleation for the given pattern. White square outlines indicate the preferred regions with weighting $w_{x,y} = 2$ during training.

The training and weighting was effective at locating pattern-specific nucleation to the preferred regions, as estimated by the Stochastic Greedy Model. Figure 1.5 show color-coded and superimposed nucleation heatmaps of $\eta_{x,y}^+$ for each training pattern on its target shape. (Individual nucleation heatmaps can be found in Section YYY, including heatmaps for nucleation on the off-target shapes.) Because there were 4 preferred regions for each shape, but 6 training patterns targeted to each shape, the strong effect of the position-dependent weights resulted in confining nucleation to the preferred regions, and thus forcing more than one shape to share the same nucleation region in some cases. Examination of how these patterns embed in the shapes (see Section 5.4), we see that although the same or overlapping nucleation regions are used, the highest-concentration tiles used by each pattern are generally different (c.f. Avogadro vs Abbott). This suggests that if all locations worked well for nucleation, there would be considerable potential for storing and recognizing more patterns per class in future improved designs that distribute nucleation throughout the shapes. (At this point we do not know whether the inferior nucleation locations were due to strand synthesis, sequence design, or other factors.)

1.6 Training and testing image sources and processing

In our work here, we also placed restrictions on the images to be recognized. First, all images used for training and testing should make use of the same total concentration, so as to control for the strong effect that absolute concentration has on nucleation – we are interested in conditions where the *pattern* of relative concentrations is the deciding factor. Second, all images should have the same histogram of pixel values, so as to control for the fact that a combination of high concentration and low concentration tiles can be more effective than uniform concentrations with the same total. This second restriction is more stringent and implies the first. Neither is necessary in principle for pattern recognition by nucleation, but eliminate some possible confounds for interpreting this first foray into the phenomenon. Expanding our understanding of pattern recognition by nucleation into the full space of possible images could reveal novel properties; for example, we might expect dilutions of a concentration pattern to be classified equivalently for a suitably slow anneal until the first nucleation occurs.

The particulars of image processing in this work are as follows. To allow reasonable mixing of reagents by the acoustic liquid handler we used, which had a discrete volume step of 25 nL, pixel values for each image were binned into 10 bins. As described in the Methods, pixel values $0 \leq p_n \leq 1$ were mapped exponentially to tile concentrations, using $c_i = c e^{3 p_n \ln 3}$, where base concentration $c = 16.67$ nM, c_i is a tile concentration and $i = \theta(n)$. Pixel value 0 corresponds to black and 1 to white. Because of pixel value binning, the pixel values were $x/9$ for integer $x \in [0, 9]$, which similarly resulted in 10 bins for tile concentrations, between 16.67 and 450 nM.

Images for the pattern recognition experiments came from a wide variety of sources. As described in Methods, each was cropped and rescaled to a 30×30 array, with grayscale levels adjusted to match a consistent histogram. We selected images that could be categorized into three classes based on the first letter of their names, with class labels and shapes ‘H’, ‘A’, and ‘M’ chosen to reflect ‘Hopfield Associative Memory’, the foundational inspiration for our perspective on multifarious self-assembly. Images were downloaded and processed in September, 2019. Quotes are from the given link or from Wikipedia.

Hodgkin. https://en.wikipedia.org/wiki/Dorothy_Hodgkin.

Public domain image of Dorothy Mary Crowfoot Hodgkin. Hodgkin “was a Nobel Prize-winning British chemist who advanced the technique of X-ray crystallography to determine the structure of biomolecules, which became essential for structural biology.” Our self-assembly work relies on the formation of geometrically ordered structures that would produce point-based diffraction patterns in the limit of large structures, much like classical homogeneous crystals. However, our structures have different DNA sequences at different sites and hence are not periodic if the sequence identity is taken into account. Our structures can be seen as a generalization of crystals to the multicomponent limit where the number of distinct species is comparable to the system size.

Hopfield. <https://alchetron.com/John-Hopfield>.

Permission for using this image obtained directly from John Joseph Hopfield. Hopfield “is an American scientist most widely known for his invention of an associative neural network in 1982.” Hopfield’s Associative Memory inspired the concepts underlying this work, even though our work exploits an inevitable physical process (nucleation) to perform pattern recognition without mimicking Hopfield neural networks element-by-element.

Horse. <https://pixabay.com/photos/horse-mold-thoroughbred-arabian-2063672>.

Pixabay’s license is “free for commercial use, with no attribution required.” The Arabian horse is “is also one of the oldest breeds, with archaeological evidence of horses in the Middle East that resemble modern Arabians dating back 4,500 years.”

Hazelnuts. <https://pixabay.com/photos/hazelnut-nuts-legume-nut-healthy-3357096>.

Pixabay’s license is “free for commercial use, with no attribution required.” As of this writing, the oldest “evidence of large-scale Mesolithic nut processing, some 8,000 years old, was found in a midden pit on the island of Colonsay in Scotland.”

Harom. <http://www.pymvpa.org/datadb/mnist.html>.

“Harom” is the word for “3” in Hungarian. The image is from the MNIST database.

H. <https://www.nist.gov/itl/products-and-services/emnist-dataset>.

The image of the letter “H” is from the EMNIST database.

Avogadro. https://en.wikipedia.org/wiki/Amedeo_Avogadro.

Public domain image of Lorenzo Romano Amedeo Carlo Avogadro. Avogadro “was an Italian scientist, most noted for his contribution to molecular theory now known as Avogadro’s law.” His law tells us how many parallel pattern recognition events ($\sim 10^{11}$) occurred in each macroscopic test tube sample (i.e. how many nuclei have formed at the time of 10% signal quenching), making us wonder how small a system could carry out the same task.

Abbott. <https://www.frontiersin.org/journals/neuroscience>.

Permission for using this image, downloaded from the Neuroscience journal Editorial Board list, was obtained directly from Laurence Frederick Abbott. Abbott is an American theoretical neuroscientist, known for his studies of neural adaptation and plasticity, including reservoir computing and competitive Hebbian learning in networks of spiking neurons. While we trained our molecular systems a la reservoir computing using a computer, our design using interaction mediating particles potentially allows learning interactions through a chemical Hebbian process in future work.

Anchovy. https://en.wikipedia.org/wiki/Anchoa_lyolepis.

Licensed under the Creative Commons Attribution 2.0 Generic license. The Dusky anchovy (*Anchoa lyolepis*) “is a species of anchovy native to the western Atlantic Ocean from New York to Brazil.”

Apples. <https://www.publicdomainpictures.net/en/view-image.php?image=17402&picture=green-apples>.

Available under the CC0 Public Domain dedication. The Granny Smith “is an apple cultivar which originated in Australia in 1868. It is named after Maria Ann Smith, who propagated the cultivar from a chance seedling.”

Aon. <http://www.pymvpa.org/datadb/mnist.html>.

“Aon” is the word for “1” in Scottish Gaelic. The image is from the MNIST database.

A. <https://www.nist.gov/itl/products-and-services/emnist-dataset>.

The image of the letter “A” is from the EMNIST database.

Mitscherlich. https://en.wikipedia.org/wiki/Eilhard_Mitscherlich.

Public domain image of Eilhard Mitscherlich. Mitscherlich “was a German chemist, who is perhaps best remembered today for his discovery of the phenomenon of crystallographic isomorphism in 1819 [...] His investigation, also in 1826, of the two crystalline modifications of sulfur threw much light on the fact that the two minerals calcite and aragonite have the same composition but different crystalline forms, a property which Mitscherlich called polymorphism.” The multifarious S+H+A+M molecular mix, capable of three distinct geometric ordered assemblies, can be seen as a generalization of the crystal polymorph concept to the limit of a large number of distinct species.

Moser. <https://www.ntnu.edu/erc-grants>.

Permission for using this image, downloaded from the Norwegian University of Science and Technology, was obtained directly from May-Britt Moser. Moser, with “her then-husband, Edvard Moser, shared half of the 2014 Nobel Prize in Physiology or Medicine, awarded for work concerning the grid cells in the entorhinal cortex, as well as several additional space-representing cell types in the same circuit that make up the positioning system in the brain.” The collective dynamics of the multifarious self-assembly model explored here is mathematically related to place cell network models that simultaneously encode multiple spatial memories through different co-localization of place fields.

Mockingbird. <https://pixabay.com/photos/polyphonic-mockingbird-2232092>.

Pixabay’s license is “free for commercial use, with no attribution required.” The northern mockingbird, commonly found in North America, “is known for its mimicking ability, as reflected by the meaning of its scientific name, ‘many-tongued thrush’”

Magnolia. https://commons.wikimedia.org/wiki/File:Magnolia_flower_Duke_campus.jpg.

Licensed under the Creative Commons Attribution 3.0 Unported license. The southern magnolia (*Magnolia grandiflora*) “is a tree of the family Magnoliaceae native to the southeastern United States.”

Mbili. <http://www.pympva.org/databdb/mnist.html>.

“Mbili” is the word for “2” in Swahili. The image is from the MNIST database.

M. <https://www.nist.gov/itl/products-and-services/emnist-dataset>.

The image of the letter “M” is from the EMNIST database.

1.7 Mixes for pattern recognition experiments

Sample mixes were designed to be possible using a particular order of automated intermediate mixes, first creating mixes of tiles at each concentration value (discretized by the pixel value bins) for each pattern, then mixing these to obtain a full sample mix for each pattern. Absolute tile concentrations were set to have a target average tile concentration of 60 nM. This resulted in slightly different discrete tile concentration bins than were used in the nucleation model. Additionally, the discrete volume steps used by the acoustic liquid handler resulted in slightly different concentrations in experiments than the target concentrations, as shown in Table 1.1. The actual concentrations (nominally) dispensed by the liquid handling robot are different from the model concentrations by no more than 3.2 %.

Table 1.1: Tile concentrations for pattern recognition.

Factor	Model (nM)	Target (nM)	Actual (nM)
1.00	16.67	17.08	16.87
1.44	24.04	24.63	24.51
2.08	34.67	35.53	35.71
3.00	50.00	51.24	51.32
4.33	72.11	73.90	73.60
6.24	104.00	106.58	107.14
9.00	150.00	153.72	153.76
12.98	216.34	221.70	223.06
18.72	312.01	319.75	321.43
27.00	450.00	461.16	464.29

References

1. Maffeo, C., Yoo, J. & Aksimentiev, A. “De novo reconstruction of DNA origami structures through atomistic molecular dynamics simulation”. *Nucleic Acids Research* **44**, 3013–3019 (2016) (cit. on p. 7).
2. Poppleton, E., Romero, R., Mallya, A., Rovigatti, L. & ulc, P. “OxDNA.org: a public webserver for coarse-grained simulations of DNA and RNA nanostructures”. *Nucleic Acids Research* **49**, W491–W498 (2021) (cit. on p. 7).
3. Fonseca, P. *et al.* “Multi-scale coarse-graining for the study of assembly pathways in DNA-brick self-assembly”. *The Journal of Chemical Physics* **148**, 134910 (Apr. 7, 2018) (cit. on pp. 7, 10).
4. Winfree, E. *Simulations of computing by self-assembly*. Tech. rep. CaltechCSTR:1998.22 (Pasadena, CA, 1998). <https://resolver.caltech.edu/CaltechCSTR:1998.22> (cit. on pp. 7, 8).
5. Evans, C. G., Hariadi, R. F. & Winfree, E. “Direct atomic force microscopy observation of DNA tile crystal growth at the single-molecule level”. *Journal of the American Chemical Society* **134**, 10485–10492 (2012) (cit. on pp. 7, 10).
6. Jiang, S., Hong, F., Hu, H., Yan, H. & Liu, Y. “Understanding the elementary Steps in DNA tile-based self-assembly”. *ACS Nano* **11**, 9370–9381 (2017) (cit. on pp. 7, 10).
7. Schulman, R. & Winfree, E. “Programmable control of nucleation for algorithmic self-assembly”. *SIAM Journal on Computing* **39**, 1581–1616 (2009) (cit. on pp. 7, 11).
8. Evans, C. G. & Winfree, E. “Physical principles for DNA tile self-assembly”. *Chemical Society Reviews* **46**, 3808–3829 (2017) (cit. on pp. 8, 9, 12).
9. Rothemund, P. W. K., Papadakis, N., Winfree, E. & Condon, A. “Algorithmic self-assembly of DNA Sierpinski triangles”. *PLoS Biology* **2**, e424 (2004) (cit. on pp. 8, 9).
10. Fujibayashi, K., Hariadi, R., Park, S. H., Winfree, E. & Murata, S. “Toward reliable algorithmic self-assembly of DNA tiles: A fixed-width cellular automaton pattern”. *Nano Letters* **8**, 1791–1797 (2008) (cit. on p. 8).
11. Gillespie, D. T. “Stochastic simulation of chemical kinetics”. *Annual Review of Physical Chemistry* **58**, 35–55 (2007) (cit. on p. 9).
12. Gibson, M. A. & Bruck, J. “Efficient exact stochastic simulation of chemical systems with many species and many channels”. *The Journal of Physical Chemistry A* **104**, 1876–1889 (2000) (cit. on p. 9).
13. Evans, C. G., Schulman, R. & Winfree, E. *The Xgrow Simulator*. 1996–2022. <https://github.com/DNA-and-Natural-Algorithms-Group/xgrow> (cit. on p. 9).
14. Schulman, R., Wright, C. & Winfree, E. “Increasing redundancy exponentially reduces error rates during algorithmic self-assembly”. *ACS Nano* **9**, 5760–5771 (2015) (cit. on p. 9).
15. Fujibayashi, K. & Murata, S. “Precise simulation model for DNA tile self-assembly”. *IEEE Transactions on Nanotechnology* **8**, 361–368 (2009) (cit. on p. 9).
16. Doye, J. P. *et al.* “Coarse-graining DNA for simulations of DNA nanotechnology”. *Physical Chemistry Chemical Physics* **15**, 20395–20414 (2013) (cit. on p. 10).
17. Woods, D. *et al.* “Diverse and robust molecular algorithms using reprogrammable DNA self-assembly”. *Nature* **567**, 366 (Mar. 2019) (cit. on p. 10).
18. SantaLucia Jr, J. & Hicks, D. “The thermodynamics of DNA structural motifs”. *Annual Review of Biophysics and Biomolecular Structure* **33**, 415–440 (2004) (cit. on p. 10).
19. Evans, C. G. *stickydesign v0.8.2*. 2019. <https://github.com/DNA-and-Natural-Algorithms-Group/stickydesign> (cit. on p. 10).
20. Bolhuis, P. G., Chandler, D., Dellago, C. & Geissler, P. L. “Transition path sampling: Throwing ropes over rough mountain passes, in the dark”. *Annual Review of Physical Chemistry* **53**. PMID: 11972010, 291–318 (2002) (cit. on p. 13).
21. Allen, R. J., Valeriani, C. & ten Wolde, P. R. “Forward flux sampling for rare event simulations”. *Journal of Physics: Condensed Matter* **21**, 463102 (2009) (cit. on p. 13).
22. Pan, A. C. & Chandler, D. “Dynamics of nucleation in the Ising model”. *The Journal of Physical Chemistry B* **108**, 19681–19686 (2004) (cit. on p. 13).

23. Oxtoby, D. W. “Homogeneous nucleation: theory and experiment”. *Journal of Physics: Condensed Matter* **4**, 7627 (1992) (cit. on p. 17).
24. Whitelam, S. & Jack, R. L. “The statistical mechanics of dynamic pathways to self-assembly”. *Annual Review of Physical Chemistry* **66**, 143–163 (2015) (cit. on p. 17).
25. Soloveichik, D. *CRN Simulator (version 2.1)*. 2022. <http://users.ece.utexas.edu/~soloveichik/crnsimulator.html> (cit. on p. 21).

Section 2

Fluorophore labels

2.1 Motivation

To assess the progress of nucleation and growth in our system, we used a combination of atomic force microscopy (AFM) and real-time fluorescence, each of which comes with strengths and limitations. The obvious strength of AFM is that we can see individual assemblies and collect statistics on their morphologies, in some cases with single-tile resolution; limitations include that it can be time-consuming to prepare samples and take images, that analysis can be hard to automate, and that deposition onto mica introduces biases: in our case, we adjusted conditions such that single tiles and small assemblies did not stick well and were washed away. For these reasons we limited AFM to end-point measurements. The obvious strength of real-time fluorescence is that it reports bulk averages as a function of time, thus providing insight into kinetics; limitations include that exactly what the fluorescence level is reporting sometimes can be hard to discern, and that modifications required to enable fluorescent readout may alter the behavior of the system. Addressing and understanding these limitations, for our experimental system, required us to explore and characterize several designs for fluorescent readout.

Due to the wide availability of real-time quantitative PCR (qPCR) machines, with their ability to precisely control temperature while measuring multicolor fluorescence in real time on parallel samples, they have become a useful tool for quantitative studies of thermodynamics and kinetics in nucleic acid biophysics and molecular programming.^{1–3} Ideally, the use of a qPCR machine would allow the quantitative measurement of the absolute or relative concentrations of free and bound tiles of multiple chosen types simultaneously throughout the course of self-assembly. Our particular aim was to simultaneously monitor four locations per shape, so as to be able to observe and distinguish both the timing of initial nucleation and the progress of further growth. However, there are several phenomena that present obstacles to converting from fluorescence measurements to absolute (or even relative) concentrations in our experimental conditions. Previously used methods for fluorescent monitoring of DNA self-assembly did not appear to be suitable for our needs. Self-assembly of 2D and 3D DNA nanostructures made of double-crossover and single-stranded tiles has been monitored using the SYBER Green I intercalating dye,^{4,5} but this method generically assesses single-stranded versus double-stranded rather than being easily targetable to specific locations within a structure. Fluorescence resonance energy transfer (FRET) between fluorophore pairs covalently attached to specific strands can provide proximity-dependent signals,^{6,7} but for multiplexing, each signal occupies more wavelength real estate than a single-fluorophore setup, and these studies monitored only a single position within the DNA object.

2.2 Initial design

Inspired by dual-labeled fluorescent probes such as TaqMan probes and molecular beacons,⁸ we initially used a single tile modified to contain a 5' fluorophore and a 3' quencher, as shown in Figure 2.1(a). When unbound, the modified strand was expected to allow free movement of the fluorophore and quencher without confining them to be near enough to cause significant quenching. When bound in a lattice, however, the two ends would be on adjacent helices and oriented so as to be near each other. This constrained position would only result from a sufficiently large structure and not, for example, from dimers: thus quenching was expected to indicate the presence of a well-formed lattice structure around the labelled tile. By modifying only a single tile, we also hoped to minimize changes to nucleation and growth of the overall structure. We refer to a tile with a fluorophore and quencher as a ‘label tile’.




Figure 2.1: An initial design for a label tile consisting of a fluorophore and quencher on a single SST. (a) As a free unstructured strand, the fluorescence will not be fully quenched. (b) To examine the label tile's behaviour, we used three patterns around a 5×5 set of tiles from shape H, surrounding ROX on tile H173 or FAM on tile H306. The 'checker' pattern had high concentrations of tiles adjacent to the label tile, while the 'half' pattern omitted a checkerboard pattern of tiles such that the full 5×5 structure could not form. An anneal and melt of the patterns around a ROX label tile and FAM label tile shows a significant *increase* in fluorescence during the anneal. (d) To examine whether DNA binding to the label tile 'stretched' the strand, reducing self-quenching, we used 'guard' strands that each bound to a third of the tile. (e) For all four fluorophores, samples with guard strands showed a significant increase in fluorescence during the anneal around 50°C .

Using this 'label tile' design in initial experiments, we found that while changes to fluorescence occurred near expected growth temperatures, the changes were not as simple as a transition from a bright unbound state, to a quenched bound state. Using 5×5 sets of tiles around label tiles to characterize their behaviour by growing small structures during annealing experiments, we found that fluorescence significantly increased when ramping from temperatures well above melting to temperatures around those favourable for growth, before quenching at lower temperatures, but only to levels slightly below those seen at high temperatures (Figure 2.1(b), 'checker' and 'uniform'). In contrast, samples missing half the tiles required for structure formation, but with adjacent tiles that could bind to the label tile, showed an increase in fluorescence as temperature decreased (Figure 2.1(b), 'half').

These results led us to suspect that, when unbound and single-stranded, the fluorophore and quencher on the label tile were both not constrained by the DNA to be close enough to quench fully, but were also not constrained to be far apart, and thus tended to interact and quench at an intermediate level. However, when some domains on the tile were bound to those on other tiles, whether in the course of nucleating a larger lattice structure or simply binding to complementary regions, the increased stiffness from the double-stranded regions did force the fluorophore and quencher apart, stopping this quenching and raising the fluorescence signal. Figure 2.1(a) shows an example intermediate structure that can occur in all the samples, but which will continue to assemble into a fully-quenched 5×5 square in all but the 'half' sample. This interpretation is consistent with the understood mechanism by which TaqMan probes, which typically have a baseline fluorescence several fold higher than molecular beacons that use an explicit hairpin stem to bring fluorophore and quencher together, enhance their fluorescence upon binding to a target.⁸

To verify this interpretation, we used 14-nucleotide 'guard strands' complementary to the label tiles as shown in Figure 2.1(c), such that when added to a sample containing only the matching label tile, they would cause the tile to be double-stranded at low temperatures. For all our label tiles, this addition resulted in higher fluorescence at temperatures lower than the guard strands' $\sim 50^\circ\text{C}$ predicted⁹ melting point, while the label tiles by themselves showed relatively little temperature dependence (Figure 2.1(d)).

While it was possible to distinguish the formation of structures through the increased and decreased fluorescence during anneals, as in 2.1(b), the complexity and inconsistency of the label tile's behaviour during nucleation motivated a second design, which was used for all subsequent experimental results.

2.3 Final design

To avoid fluorophore-quencher interactions outside of lattice formation, our second design, shown in Figure 2.2(a), placed the fluorophore and quencher on two separate tiles - a ‘label pair’ - with no complementary regions. In addition to avoiding the intra-tile interactions of the previous design, the two tiles in a label pair cannot form a dimer that might exhibit quenching, even at relatively low temperature. While normal SSTs in this configuration would not have terminal ends close enough for contact quenching, in this design, one of the two tiles had its orientation reversed, with a crossover in a lattice on the opposite side; as a result, in a well-formed lattice, the fluorophore and quencher are located on facing sides of adjacent helices. The same tile, when used without a fluorophore, does not use the reversed orientation.

Using this label design, formation of structures caused a monotonic corresponding decrease from an unquenched, high fluorescence, to a quenched, low fluorescence signal, subject to the temperature dependence of the fluorophore itself (data not shown). This behaviour was seen with reasonable consistency in preliminary multifarious growth experiments, and could be compared with AFM images of the resulting structures.

Given these results, we might hope to construct a formula f linking fluorescence to assembled (as opposed to free) tile concentration, e.g. $c_i^{\text{assem}} = f(c_i, F, T)$, for a particular fluorescent label tile i , fluorescence reading F , temperature T , and total tile concentration c_i . Doing so would be possible using suitable calibration experiments if, for example, free tiles and assembled tiles each had well-defined fluorescence levels at each temperature, independent of other DNA molecules in solution. (Here, we take ‘assembled tiles’ to mean those within assemblies of at least four strands that adopt the lattice conformation that brings fluorophore and quencher close to each other, while by an abuse of terminology we take ‘free tiles’ to mean those that are free or bound to others in locally linear or tree-like chains that don’t fold the fluorophore and quencher tiles as in the lattice. It would be the concentration of this population that we could infer.)

In initial control experiments (Figure 2.2(c)), we found that strands that bind directly to the fluorophore tile substantially increase its fluorescence, as expected. But more surprisingly, this effect only moderately decreases at higher temperatures where the binding is predicted to be unfavorable, as well as in the presence of unrelated DNA with little expected binding. To characterize this behaviour further, we took melting curves of a 5×5 square of 50 nM tiles around the ROX label pair in H, along with a variable amount of an equal mixture of tiles unique to H, outside a 7×7 region around the same label pair (Figure 2.2(b)). Owing to the roughly checkerboard pattern of shared and unique tiles, this unique tile mix approximated a collection of unrelated, non-interacting DNA, which was not complementary to either the tiles in the 5×5 target assembly, or to other strands within the mixture.

As shown in Figure 2.2(d), increasing the total concentration of unrelated, non-specific DNA (i.e., $\sum_i c_i$ for strand concentrations c_i) in a sample increased fluorescence at all temperatures, in a seemingly consistent way unaffected by the quenching when the 5×5 region grew and melted. The effect was present even when all assemblies were clearly melted, above 65 °C, and appears to be roughly linear between 0 and 20 tM.

Knowing that in the absence of Tween, PEG, carrier DNA, or the equivalent (which we did not use in these experiments), low concentrations of DNA can partially absorb onto the walls of test tubes and pipette tips, we explored the hypothesis that this was related to our observations – that somehow the excess strands were playing the role of carrier DNA. While we used coated tubes designed to minimize DNA adsorption (0.5 mL DNA LoBind tubes, Eppendorf), and pipette tips (Eppendorf LoRetention), we considered whether fluorescently-labeled DNA (and assembled structures containing the label pairs) could have been absorbed to the walls when excess strands were not present, thus yielding low fluorescence measurements, but released from the walls when excess strands were available to replace them, thus increasing the measured fluorescence. Quantitatively, the observed effect seemed too large for this mechanism, and further investigation also ruled it out as a significant factor.

However, strands with regions complementary to regions on the fluorophore tile also appear to affect fluorescence, to a greater extent than non-specific DNA, and this effect also persists even at temperatures where no binding would be expected to be stable.

To measure this effect, we took melting curves of samples with all tiles for constructing H at 50 nM, along with a 10x concentration checkerboard flag pattern, either surrounding the label pair, or in a position far from the label pair. Both types of samples had the same total DNA concentration, but the sample with the label pair inside the high-concentration pattern, thus having high concentration of tiles adjacent to the fluorophore and quencher, had a significantly higher concentration of strands with regions complementary to regions on the fluorophore and quencher strands. In an anneal and melt, shown in Figure 2.2(e), the sample with high concentration adjacent strands had a significantly higher fluorescence above the melting temperature of the structure, but when structures had formed, had similar fluorescence. The raw fluorescence level at low temperatures, regardless of whether the excess tiles were adjacent or not, was consistent with that of the sample in Figure 2.2(d) that had the most similar




Figure 2.2: A two-tile label pair, separating the fluorophore and quencher, gives quenching from lattice growth without significant interference from intermediate states, but effects on fluorescence related to other strands complicate quantitative interpretation of fluorescence in terms of absolute tile and structure concentrations. (a) The label scheme uses two nearby tiles, along the same helices but not binding to each other directly, with one tile reversed in orientation, resulting in 5' modifications on the two tiles being constrained to be near to each other in a well-formed lattice, but not constrained in any potential complex of two or three tiles. (b) To characterize the label pair behaviour, we ran experiments with several combinations of components, based around a ROX fluorophore on tile H173 and a 5×5 surrounding area. A ‘filler’ mix included all tiles unique to H outside a 7×7 region around the fluorophore tile, intended to provide a mix of non-interacting tile-like DNA strands. Complete H structure samples used the same ROX fluorophore tile, with its adjacent quencher tile, and 5×5 area, but with one of two possible checkerboard patterns (blue or orange) of increased concentrations. (c) An anneal and melt with the (H173 with ROX) fluorophore tile and no quencher tile, including only the fluorophore tile, the fluorophore tile with $10 \mu\text{M}$ total filler concentration, and the fluorophore tile with each of the four tiles adjacent to it that each bind by one domain. We interpret the anomalously low fluorescence at the beginning of the anneal, which started by cooling from just below 60°C , to the samples not yet having established equilibrium. (d) An anneal and melt of a 5×5 block with varying total concentration of filler tiles. The filler strands are expected to remain unbound at all temperatures. (e) An anneal and melt of a complete H structure, including the same label pair. At high temperature, the blue and orange samples are expected to have the same total concentration of strands, but different concentrations of tiles adjacent to the label. At low temperature, if structures have completely grown, the two samples are expected to have the same concentration of structures and of excess, unbound strands. For c-e, samples were held at 70°C for 10 minutes, ramped to 57°C at $1^\circ\text{C}/\text{min}$, held at 57°C for 20 minutes, ramped from 57°C to 35°C at 6 minutes per 0.2°C , then ramped from 35°C to 67°C at 4 minutes per 0.2°C . (Note that the extent of hysteresis depends on ramp speed and reflects the nucleation barrier kinetics and melting kinetics.)

amount of excess DNA, suggesting a lack of sequence-specificity for fluorophores within assembled structures. This also rules out the possibility that the fluorescence increase was just due to unintentional excess of fluorophore tiles over non-labeled tiles (noting that label pair tiles were ordered purified, while all other tiles we used unpurified).

Our conclusions are as follows:

1. The mostly-quenched fluorescence levels of fluorophores within assembled structures, with a nearby quencher, increase similarly in the presence of unrelated or related single-stranded DNA in solution.
2. The mostly-unquenched fluorescence levels of fluorophores on free strands increase in the presence of unrelated single-stranded DNA, with further increases in the presence of related single-stranded DNA that has some sequence complementarity.
3. The free fluorophore tile's temperature dependence is small by comparison.
4. Accounting for these effects to obtain quantitative inference of tile and assembly concentrations from fluorescence data would be fraught.
5. In these experiments, we see that time/temperature of the formation and melting transitions is not very sensitive to the above factors influencing absolute fluorescence.

It seems likely that using FRET instead of direct fluorescence would alleviate some of these issues, at the cost of having fewer label pair tiles per sample. However, given the surprising influence of excess DNA even for fluorophore/quencher pairs within the assembled lattice context, it is not a foregone conclusion that FRET eliminate these confounds entirely.

As a separate matter, it is well-known that incorporation of fluorophores and quenchers can affect the thermodynamics of DNA hybridization and self-assembly reactions,¹ often being beneficial on the order of up to 2 kcal/mol. This is especially a concern for us, if the modified tiles are part of a critical nucleus that controls nucleation rates. Consistent with this, although also possibly influenced by the reverse orientation¹⁰ of the fluorophore-labeled tile, the AFM-based counts for samples containing label pair tiles near the center of the nucleation region were often notably higher than those that used label tiles elsewhere but otherwise had the same tile concentration pattern (see the table in Section 5.3.1).

References

1. You, Y., Tataurov, A. V. & Owczarzy, R. “Measuring thermodynamic details of DNA hybridization using fluorescence”. *Biopolymers* **95**, 472–486 (2011) (cit. on pp. 29, 33).
2. Padirac, A., Fujii, T. & Rondelez, Y. “Quencher-free multiplexed monitoring of DNA reaction circuits”. *Nucleic Acids Research* **40**, e118–e118 (2012) (cit. on p. 29).
3. Cherry, K. M. & Qian, L. “Scaling up molecular pattern recognition with DNA-based winner-take-all neural networks”. *Nature* **559**, 370–376 (2018) (cit. on p. 29).
4. Wang, W. et al. “Self-assembly of fully addressable DNA nanostructures from double crossover tiles”. *Nucleic acids research* **44**, 7989–7996 (2016) (cit. on p. 29).
5. Sajfutdinow, M., Jacobs, W. M., Reinhardt, A., Schneider, C. & Smith, D. M. “Direct observation and rational design of nucleation behavior in addressable self-assembly”. *Proceedings of the National Academy of Sciences* **115**, E5877–E5886 (2018) (cit. on p. 29).
6. Jiang, S., Hong, F., Hu, H., Yan, H. & Liu, Y. “Understanding the elementary Steps in DNA tile-based self-assembly”. *ACS Nano* **11**, 9370–9381 (2017) (cit. on p. 29).
7. Ke, Y. et al. “DNA brick crystals with prescribed depths”. *Nature Chemistry* **6**, 994–1002 (2014) (cit. on p. 29).
8. Huang, Q. et al. “Multiplex fluorescence melting curve analysis for mutation detection with dual-labeled, self-quenched probes”. *PloS ONE* **6**, e19206 (2011) (cit. on pp. 29, 30).
9. Zadeh, J. N. et al. “NUPACK: Analysis and design of nucleic acid systems”. *Journal of Computational Chemistry* **32**, 170–173 (2011) (cit. on p. 30).
10. Wei, B. et al. “Design space for complex DNA structures”. *Journal of the American Chemical Society* **135**, 18080–18088 (2013) (cit. on p. 33).

Section 3

Tile sequences and layout

3.1 Sequences

Tile names in this table are of the form [LETTER][TILE NUMBER], where the tile number starts from 0 (and does not depend on the letter), and the letter denotes what shapes the tile appears in. S tiles appear in all three shapes, H/A/M tiles appear only in the corresponding shape, and I/B/N tiles appear in two shapes: A+M, H+M, and H+A, respectively, akin to the IUPAC naming for degenerate DNA bases, where the letter is the letter that follows the excluded shape in the alphabet. Spaces in sequences correspond to divisions between glues. Glues are listed from 5' to 3'. They do not indicate complementarity: as we do not consider tile rotations, a glue in position 1 will bind to 3, and position 2 to position 4. Glues with the same names in these two positions (1/3 or 2/4) will be reverse complements of each other: for example, glue 3 of H1 is the reverse complement of glue 1 of H0. The exception to this is for positions that are “null” glues, which are simply 10nt or 11nt all-T regions, on tiles that will form the edges of shapes. On diagrams where the H/A/M shapes are upright, such as in Section 3.3, the glue positions in these sequences will correspond with East, North, West, and South directions, respectively. Null glue names refer to the length in nucleotides (as tiles alternate between having 10/11/11/10 nt domains and 11/10/10/11 nt domains), and, for consistency reasons, a rotated cardinal direction.

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
H0	TGGGAATATCT TTTTTTTTTT TTTTTTTTTT TTCACTGTTGTT	e1	nullT10W	nullT10N	e2
S1	AACTTCTCGA TTTTTTTTTT AGATATTCCCA ATCACCCAAA	e3	nullT11W	e1	e4
N2	AGGTAAAGAGA TTTTTTTTTT TCGAGAAAGTT TAAAAGGCCAA	e5	nullT10W	e3	e6
S3	TCACATACGT TTTTTTTTTT TCTCTTAACCT AATCGATCCA	e7	nullT11W	e5	e8
H4	TTAGTATCGGT TTTTTTTTTT ACGTATGTGA TACATGATGGA	e9	nullT10W	e7	e10
N5	ACTGCTTCCTT TTTTTTTTTT ACCGATACTAA AATCGTTCCCT	e11	nullT11W	e9	e12
H6	TTGTTCATTGT TTTTTTTTTT AAGAACAGT TGGTGAGAGTTA	e13	nullT10W	e11	e14
H7	TTTTTTTTT TTTTTTTTTT ACAATGAACAA TTCTCGACAT	nullT10S	nullT11W	e13	e15
H8	TACGGTATTGA TTTTTTTTTT TTTTTTTTTT AGGAGTTTACT	e16	nullT10W	nullT10N	e17
S9	TAGCATCACA TTTTTTTTTT TCAATACCGTA ACTAGCACAA	e18	nullT11W	e16	e19
N10	AAGATACGAGA TTTTTTTTTT TGTGATGCTA TTTCATAGGGA	e20	nullT10W	e18	e21
S11	TTTCGCTCTA TTTTTTTTTT TCTCGTATCTT AAAGAACCCA	e22	nullT11W	e20	e23
H12	AAGATACGAGA TTTTTTTTTT TAGAGCGAAA TGTGATAGACA	e20	nullT10W	e22	e25
S13	AACGTTACCT TTTTTTTTTT TCTCGTATCTT AGCCTTCTT	e26	nullT11W	e20	e27
H14	TTTCGGATAGA TTTTTTTTTT AGGTAAACGTT TTAGTAGGTGT	e28	nullT10W	e26	e29
S15	TTTTTTTTT TTTTTTTTTT TCTATCCGAAA ACTCAACACA	nullT10S	nullT11W	e28	e30
S16	TTACACCGAT AACACATGAA TTTTTTTTTT TCACCAAACCT	e31	e2	nullT11N	e32
B17	TGCGATATTTT TTTGGGTGAT ATCGGTGAA AGCTAGTGTAA	e33	e4	e31	e34
N18	TTTCCCCTGA TTTGCCTTTA AAAATATCGCA TATCAACGCT	e35	e6	e33	e36
H19	AAGGGATTAGT TGGATCGATT TCAGGGAAAA ATTGTTGACA	e37	e8	e35	e38
S20	AAACTCCGAA TCCATCATGTA ACTAATCCCTT TTGTACCACA	e39	e10	e37	e40
N21	TGGATGATACT AGGAACGATT TTCGGAGTTT TCAAGAAGGTA	e41	e12	e39	e42
N22	TCTTTCACGT TAATCTCACCA AGTATCATCCA AAAACGCAAT	e43	e14	e41	e44
H23	TTTTTTTTTT ATGTCGAGAA ACGTGAAGA TGAGGTATGAT	nullT11S	e15	e43	e45
S24	ACAGCCATTA AGTAAACTCCT TTTTTTTTTT TTCAACTCGT	e46	e17	nullT11N	e47

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
B25	TGCAGTTTTA TTGTGCTAGT TAATGGCTGT ATTAGTGGGTA	e48	e19	e46	e49
N26	TCATCGCTA TCCCTATGAAA TAAAAACTGCA AGTCAAACCT	e50	e21	e48	e51
H27	TGGCAAAATAA TGGGTTCTTT TAGCGATGAA AGTATTACGGA	e52	e23	e50	e53
B28	ACTTGTTCCT TGTCTATCACA TTATTTGCCA ACCACTTCTT	e54	e25	e52	e55
H29	AGTAGTGAAGT AAGAAAGGCT AGGAACAACT ATAGACAGGAA	e56	e27	e54	e57
S30	TTCTACGCAT ACACCTACTAA ACTTCACTACT ATCACCCCAA	e58	e29	e56	e59
H31	TTTTTTTTTT TGTGTTGAGT ATGCGTAGAA TAGGAAGACAA	nullT11S	e30	e58	e60
B32	TTGAGGGATAT AGTTGGTGA TTTTTTTTT TCGAGAGATAA	e61	e32	nullT10N	e62
S33	TTCGTACACA TAACACTAGCT ATATCCCTCAA TGCTCATCAT	e63	e34	e61	e64
B34	TGGCTATTAGA AGCGTTGATA TGTGTACGAA ACGGAAATTTT	e65	e36	e63	e66
S35	ACCTCACATT TGTCAAACAAAT TCTAATAGCCA ACCCATTTC	e67	e38	e65	e68
H36	ATGGGATTAGA TGTGGTACAA AATGTGAGGT TTTTAAGCGT	e69	e40	e67	e70
S37	ACCCCTCAAT TACCTCTTGA TCTAACCCCCAT AACACCAGT	e71	e42	e69	e72
B38	AGTAGACTGAA ATTGCGTTT ATTGAAGGGT TAATACGAGGA	e73	e44	e71	e74
S39	TTTTTTTTT ATCATAACCTCA TTCAGTCTACT TATGCACTCA	nullT10S	e45	e73	e75
H40	TGAGGACTATT ACGAGTTGAA TTTTTTTTT TCGAGAGATAA	e76	e47	nullT10N	e62
S41	TGCATATCCA TACCCACTAAT AATAGTCTCA TTAGCAACCA	e78	e49	e76	e79
H42	TGAGTTTTTA AGGTTTGACT TGGATATGCA AAAATTAGCGT	e48	e51	e78	e81
S43	TTGACACCAT TCCGTAATACT TAAAAACTGCA TTACCCAACA	e82	e53	e48	e83
H44	AAGTTAGGGTA AAGAAAGTGGT ATGGTGTCAA TGGTATTTCAGA	e84	e55	e82	e85
S45	ACCCCTCTTT TTCCTGTCTAT TACCCCTAACCTT ATTGCCACTA	e86	e57	e84	e87
N46	AGGGATGTAAT TTGTGGTGAAT AAAGAAAGGGT ATATGGTAGCT	e88	e59	e86	e89
S47	TTTTTTTTT TTGTCTTCCTA ATTACATCCCT ATTACCCGTA	nullT10S	e60	e88	e90
S48	ATGTCCACTT TTATCTCTGA TTTTTTTTTT TCGATCCAAT	e91	e62	nullT11N	e92
B49	AATAGGGTACA ATGATGAGCA AAGTGGACAT AGGATATAGCA	e93	e64	e91	e94
S50	ATTGACCCCTT AAAATTTCCGT TGTACCCCTATT ATTACAGCCA	e95	e66	e93	e96
H51	TGGTAGTAACT TGAAATGGGT AAGGGTCAAT ATAGAGTTGGA	e97	e68	e95	e98
S52	AACTTCAGCT ACGCTTAAAAA AGTTACTACCA ATCACGCTTA	e99	e70	e97	e100
H53	TGGAAGAGTAT ACTGGTGAATT AGCTGAAGTT AAATTGGAACA	e101	e72	e99	e102
S54	AAACCACTCT TCCTCGTATTA ATACTCTTCCA ATTACTGCCA	e103	e74	e101	e104
H55	TTTTTTTTTT TGAGTGCATA AGACTGGTTT ATTTTGACGT	nullT11S	e75	e103	e105
S56	TTACCGACAT TTATCTCTGA TTTTTTTTTT ATCCATCGAA	e106	e62	nullT11N	e107
H57	AGAATTAGGGA TGGTTGCTAA ATGTCGGTAA TAGAGAATCGA	e108	e79	e106	e109
S58	ACTGAACCTT ACGCTAATTTC TCCCTAATTCT ACTCGTAACA	e110	e81	e108	e111
H59	ATCTGGAGTA TGTTGGTAA AAGGTCAGT TTTTGGTTCAA	e112	e83	e110	e113
B60	ACCCCTACAAA TCTGAATACCA TACTCCAAGAT AATCCTTGCT	e114	e85	e112	e115
B61	ACAAGGATAGA TAGTGGCAAT TTGTGAGGCT TAGGGTAACAT	e116	e87	e114	e117
N62	AAGTCACCAA AGCTACCATAT TCTATCCTTGT TTCTTCACGT	e118	e89	e116	e119
H63	TTTTTTTTT TACGGTGAAT TTGGTACTT ATGCAAAGATT	nullT11S	e90	e118	e120
H64	AAGTACTGAGA ATTGGATCGA TTTTTTTTTT AGTAGGAAGAA	e121	e92	nullT10N	e122
S65	ACTGCCATAA TGCTATATCCT TCTCAGTACTT ATGCATACCA	e123	e94	e121	e124
H66	TTATCTGTGGA TGGCTGTAAAT TTATGGCAGT AGACTTAGGAA	e125	e96	e123	e126
S67	TACGTTCCAA TCCAACCTAT TCCACAGATAA ACATCTGCTT	e127	e98	e125	e128
H68	ATACTGGATGA TAAGCGTGA TTGGAACGTA AATAGTCTGGT	e129	e100	e127	e130
S69	AACCTCGAAA TGTTCCAATTTC CATCCAGTAT AGCCTTTCTT	e131	e102	e129	e27
H70	TGTTGAACCTT TGGCAGTAAT TTTCGAGGTT AGCTAGTGTAA	e133	e104	e131	e34
B71	TTTTTTTTT ACGTAAAAAA AAGTTCAAACCA ATCTCTGCA	nullT10S	e105	e133	e135
H72	AGAACTAAGGA TTCGATGGAT TTTTTTTTTT AACTATGAGGA	e136	e107	nullT10N	e137
S73	TAGCCAACAA TCGATTCTCTA TCCTTAGTTCT TCGACTTCTT	e138	e109	e136	e139
H74	ACGAGATAGAA TGTTACGAGT TTGTGGCTA AATGTGGAAAA	e140	e111	e138	e141
N75	AACGTCTTCT TTGAACCAAAA TTCTATCTCGT TCCAATCCAA	e142	e113	e140	e143
H76	AGTGATTAGGA AGCAAGGATT AGAAGACGTT AGGTTCTAGTT	e144	e115	e142	e145
S77	AGCTTCATCA ATGTTACCCCTA TCCTAATCACT ATGCACTACT	e146	e117	e144	e147
H78	TGGACAAATTACGTGAAGAA TGATGAAGCT ATGGGATCATA	e148	e119	e146	e149
B79	TTTTTTTTT AATCTTCTGAT AAATTTGTCCA TAACACCCCAA	nullT10S	e120	e148	e150
S80	ACATCCATGA TTCTTCCTACT TTTTTTTTTT ATTACCCCTT	e151	e122	nullT11N	e152
H81	TAGAGGAAAGT TGGTATGCAT TCATGGATGTT AGAACTATGGT	e153	e124	e151	e154
S82	AACCCCTCTT TTCCCTAAGTCT ACTTTCTCTCA ACTAACGACA	e155	e126	e153	e156
H83	ACAGTAGGATT AAGCAGATGT AAAGAGGGTT ATTATCAGGGT	e157	e128	e155	e158

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
N84	AATCACAGCT ACCAGACTATT AATCCTACTGT TCCGATCAAA	e159	e130	e157	e160
N85	AGGGTTCTAAT AAGAAAGGCT AGCTGTGATT AATAGTCTGGT	e161	e27	e159	e130
S86	TTCGCATCTA TAACACTAGCT ATTAGAACCCCT TCCGATCAAA	e163	e34	e161	e160
B87	TTTTTTTTTT TGCGAGAGAT TAGATCGGAA AGTGAACAAAA	nullT11S	e135	e163	e165
S88	ACACAGAACCA TCCTCATAGTT TTTTTTTTTT TCATAGCCAA	e166	e137	nullT11N	e167
H89	AGAGTTTAGGA AAGAAAGTCGA TGTCTGTGT AGATCAGTGT	e168	e139	e166	e169
S90	AGCTTCATCA TTTTCCACATT TCCTAAACTCT TAGCAAACCA	e146	e141	e168	e171
B91	TGGAGATACAT TTGGATTGGA TGATGAAGCT TCTAAGTGTGA	e172	e143	e146	e173
S92	AAACCCAGAT AACTAGAACCT ATGTATCTCCA ATCACCCAAA	e174	e145	e172	e4
H93	TATCGGAAGTA AGTAGTGCA ATCTGGGTTT AGCAAAAGAAA	e176	e147	e174	e177
N94	TGCTCCATTA TATGATCCCATT TACTTCGATA AAACCTCCCT	e178	e149	e176	e179
N95	TTTTTTTTTT TTGGGTGTTA TAATGGAGCA ACTTTAGAGGT	nullT11S	e150	e178	e180
H96	AGCTTAGATGA AAGGGTGAAT TTTTTTTTTT AGAGCTATGAT	e181	e152	nullT10N	e182
B97	TCTTGCTCT ACCATAGTTCT TCATCTAAGCT TTCTGACCAA	e183	e154	e181	e184
H98	TTAGGATGACA TGTCGTTAGT AGAACGAAGA TGGAGTAAGAT	e185	e156	e183	e186
B99	TCCCCAAAGA ACCCTGATAAT TGTCATCCTAA ATCGAACCTT	e187	e158	e185	e188
B100	AGTTGATAGGT TTTGATCGGA TCTTTGGAA TTTAAATGCGA	e189	e160	e187	e190
B101	TTTACACCGA ACCAGACTATT ACCTATCAACT TTCGACACTT	e191	e130	e189	e192
H102	TATATGGTCGT TTTGATCGGA TCGGTGAAA AGAGTTAGGAA	e193	e160	e191	e194
B103	TGACAACCAT TTTGTTCACT ACGACCATAA TCATTGCCATA	e195	e165	e193	e196
H104	AGATGTGATCT TTTTTTTTTT ATGGTTGTCA ATGATTGCAAT	e197	nullT10W	e195	e198
S105	ACCTCACATT TTTTTTTTTT AGATCACATCT AACTCCGATT	e67	nullT11W	e197	e200
H106	AGTTAAGAGA TTTTTTTTTT AATGTGAGGT AATTTTCGGT	e5	nullT10W	e67	e202
N107	ACCAGCAATA TTTTTTTTTT TCTCTAACCT ACACCATCAT	e203	nullT11W	e5	e204
H108	TTTCGGATAGA TTTTTTTTTT TATTGCTGGT AAAAGTTGTGT	e28	nullT10W	e203	e206
S109	TTTCCCTTGA TTTTTTTTTT TCTATCCGAAA TCATTGCCATA	e207	nullT11W	e28	e196
H110	AGTTTGTGTTGT TTTTTTTTTT TCAAGGGAAA TATATGACGGA	e209	nullT10W	e207	e210
S111	AAACTTCCA TTTTTTTTTT ACAAAACAAACT ATCACCCACAA	e211	nullT11W	e209	e59
H112	TTCTATGTGGA TTGGCTATGA TGGGAAGTTT AGATTATCGGA	e213	e167	e211	e214
S113	ATCCACCATT TACACTGATCT TCCACATAGAA TCATTGCTCT	e215	e169	e213	e216
H114	TCGTTTAGAGA TGGTTTGCTA AATGGTGGAT TCTTAGAAGGT	e217	e171	e215	e218
N115	AACCCCTCTT TCACACTTAGA TCTCTAACGA ACTGTCACAT	e155	e173	e217	e220
N116	AGAACTAAGGA TTTGGGTGAT AAAGAGGGTT AAAGAGAGGAA	e136	e4	e155	e222
S117	ATATGCACCA TTTCTTTTGCT TCCTTAGTTCT ATGACCCAAT	e223	e177	e136	e224
H118	AAGTCGAGTAT AAGGAGGTT TGGTGCATAT AGAGTAGTGT	e225	e179	e223	e226
B119	TTTTTTTTT ACCTCTAAAGT ATACTCGACTT TGAACCAACT	nullT10S	e180	e225	e227
S120	ACTGAACCTT ATCATAGCTCT TTTTTTTTTT TCTACCTCCT	e110	e182	nullT11N	e229
N121	TGGAGATACAT TTGGTCAGAA AAGGTTCACTG TAAGAAGGTCA	e172	e184	e110	e231
B122	ACTCGCAATA ATCTTACTCCA ATGTATCTCCA ATTGCCACTA	e232	e186	e172	e87
H123	AGAACAGAGTA AAGGTTCGAT TATTGCGAGT TCTGGGATTAT	e234	e188	e232	e235
S124	TTGACATCCA TCGCATTAAA TACTCTGTTCT ACTTGCCTAA	e236	e190	e234	e237
H125	TTGAGAGTAGT AAGTGTGCAA TGGATGTCAA ACAATGGAAAA	e238	e192	e236	e239
S126	TCGATTCCCTT TTCCCTAACTCT ACTACTCTCAA ACTGCTACAT	e240	e194	e238	e241
H127	TGGATGATACT TAGGCAATGA AAGGAATCGA TTCGGTATAGA	e41	e196	e240	e243
S128	ACGTCCATAA ATTGCAATCAT AGTATCATCCA ATTGCCACTA	e244	e198	e41	e87
H129	TGGTGTGATT AATCGGAGTT TTATGGACGT AGAGTTAGGAA	e246	e200	e244	e194
S130	AATCCGAACCT ACCGAAAAATT AATCAACAAACA TGCTCATCAT	e248	e202	e246	e64
N131	ACTGAGGATAA ATGATGGGT AGTCGGATT TAATAGTCGGA	e250	e204	e248	e251
S132	TTCCCAAAGA ACACAACTTTT TTATCCTCAGT TCAAGAACCA	e187	e206	e250	e253
B133	ATCTTGGAGTA TAGGCAATGA TCTTTGGGAA TATAGACGTGT	e112	e196	e187	e255
N134	ACCTGACATT TCCGTATATA TACTCCTAACAGAT AGAAAACACCT	e256	e210	e112	e257
B135	AACGTGAAATT TTGTGGTGAT AATGTCAGGT ATGTAAGTAGCA	e258	e59	e256	e259
S136	AATCAGCCCT TCCGATAATCT AATTTCACGTT ACTTGCCTAA	e260	e214	e258	e237
H137	TAGTCAGTTGA AGAGCAATGA AAGGCTGATT TTAGTCAGGA	e262	e216	e260	e263
S138	ACTGAACCTT ACCTTCTAAAGA TCAACTGACTA TTTCGTCTT	e110	e218	e262	e265
H139	TGTAGATGAGT ATGTGACAGT AAGGTTCACTG TAGGGATGAAA	e266	e220	e110	e267
N140	ATCGTAACCA TTCCCTCTCTT ACTCATCTACA ATCTCAGCAT	e268	e222	e266	e269
N141	AGGAAGGAATA ATTGGGTCACT TGGTTACGAT TGAAGTAAGGA	e270	e224	e268	e271
S142	TCTTCACGA AACACTACTCT TATTCCCTCCT AGCTCTCATT	e272	e226	e270	e273

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
H143	TTTTTTTTTT AGTTGGTCA TCGTGAAGA ATTGAAGCAAA	nullT11S	e227	e272	e274
H144	AGAGATGACTT AGGAGGTA AAGTCATCTCT TTTCCGTCTT	e275	e229	nullT10N	e276
S145	TCAACCCATT TGACCTTCTTA AAAGTCATCTCT TTTCCGTCTT	e277	e231	e275	e265
H146	ACTTGTAGGAT TAGTGGCAAT AAATGGGTGA AGGAAAGACTA	e279	e87	e277	e280
S147	TAACTGCACA ATAATCCAGA ATCCTACAAGT TCACAACGTA	e281	e235	e279	e282
H148	ATGTGAGTAGT TTAGGCAAGT TGTGCGAGTT AAATTGGAACA	e283	e237	e281	e102
S149	ATATCGCACA TTTTCCATTGT ACTACTCAT AATCTCGACA	e285	e239	e283	e286
H150	TGTATGAAGGA ATGTAGCGAT TGTGCGATAT AAAGGCTAGTA	e287	e241	e285	e288
N151	ACACACAAGA TCTATAACGAA TCCCTCATACA AGTCACACAA	e289	e243	e287	e290
N152	AGAGTTTAGGA TAGTGGCAAT TCTTGTGTG TAGAGAACATCGA	e168	e87	e289	e109
S153	TGTACCCAAA TTCCTAACTCT TCCTAAACTCT TCATGCCCTA	e293	e194	e168	e294
H154	ATGTAGGACTT ATGATGAGCA TTTGGGTACA TGGAGACTAAT	e295	e64	e293	e296
N155	ACATGTACCA TCCGACTATTAA AAGTCCTACAT TCATCACCAT	e297	e251	e295	e298
H156	ATGGATGTCATA TGGGTTCTGA TGTGACATGT TAATAGTCGGA	e299	e253	e297	e251
S157	TTCCAACAGA ACACGTCTATA TAGACATCCAT TTCAACCCAT	e301	e255	e299	e302
H158	ATGTAGGGTAA AGGTGTTCT TCTGTTGAA AAGAGAGTGTAA	e303	e257	e301	e304
S159	ACCCTTCTTT TGCTACTACAT TTACCCCTACAT ACCATCCAAA	e86	e259	e303	e306
H160	TAGAAGGCATA TTAGGCAAGT AAAGAAGGGT AGCAAAGAAAT	e307	e237	e86	e308
S161	TTCGCTCTAA TCCTGAACCTAA TATGCCCTCTA AGCACTTACA	e309	e263	e307	e310
H162	TGGATAGCTTA AAGACGGAA TTAGAGCGAA ATTAGTGGGTA	e311	e265	e309	e49
B163	ACGATCATCA TTTCATCCCTA TAAGCTATCCA TTCCGTACAA	e313	e267	e311	e314
H164	AGGTATCGTAT ATGCTGAGAT TGATGATCGT AAAATGCAAGA	e315	e269	e313	e316
N165	ACGATCATCA TCCTTACTTCA ATACGATACCT TCAAGACACA	e313	e271	e315	e318
S166	TAAAATTGGCA AATGAGAGCT TGATGATCGT AATGTCAGGA	e319	e273	e313	e320
S167	TTTTTTTTTT TTTGCTTCAAT TGCCAAATTAA TCAATCAGCT	nullT10S	e274	e319	e321
B168	AACGTCTTCT ACGACTAACTA TTTTTTTTTT TCACCGTTAT	e142	e276	nullT11N	e323
N169	TTAGGTAGCAT AAGACGGAA AGAACAGCTT ATAGTGAGGAA	e324	e265	e142	e325
N170	ACAACTTCGA TAGTCTTCTCT ATGCTACCTAA TTACCAACCAA	e326	e280	e324	e327
H171	AATGCAGATT TACGTTGTGA TCGAAGTTGT TTTCATTTGGT	e328	e282	e326	e329
S172	ACACTCCAAA TGTCCAATTAA AATCTGCATT ATTGTCCCTT	e330	e102	e328	e331
H173	TTTGATTCTG TGTGAGATT TTTGGAGTGT ACTTTAGAGGT	e332	e286	e330	e180
S174	ACCATAACGA TACTAGCCTT ACGAACATAAAA AGCATCAACT	e334	e288	e332	e335
H175	AGGTATCGTAT TGTGTGAAC TCGTTATGGT TCAAGAACGTA	e315	e290	e334	e42
S176	ACTTGTCTCT ACGACTAACTA ATACGATACCT AAATCCTGCT	e54	e109	e315	e339
B177	TTAGGTAGCAT TAAGGCATGA AGGAACAAAGT AGGTTAGACAT	e324	e294	e54	e341
S178	AAACCTGCTA ATTAGTCTCCA ATGCTACCTAA ATTACGCTCT	e342	e296	e324	e343
B179	TAGGAACAGAA ATGGTGATGA TAGCAGGTTT ACAGGAAGATA	e344	e298	e342	e345
S180	AATCACACGT TCCGACTATTAA TTCTGTCTCTA AGCCTCATTT	e159	e251	e344	e347
H181	TGGTAAGTACT ATGGGTTGAA AGCTGTGATT AATGTCAGGA	e348	e302	e159	e320
S182	TACGTTCCAA TACACTCTCTT AGTACTTACCA TCGACTTTCA	e127	e304	e348	e351
N183	AGAAGTCAGAT TTTGGATGTT TTGGAACGTA AGCAAAGAAA	e352	e306	e127	e177
N184	ATACACACGT ATTTCTTGCT ATCTGACTTCT AGCTCTCATT	e354	e308	e352	e273
H185	ATGGTTTTGTT TGTAAGTGTCT ACGTGTGTAT TGAAAGAGGT	e356	e310	e354	e357
S186	ACGTATCACAA TACCCACTAAT AACAAAACCAT ACCAACGCTAA	e358	e49	e356	e359
H187	TTGAGGGATAT TTGTCAGGAA TGTGATACGT ATATGGTAGCT	e61	e314	e358	e89
S188	ACACACAAGA TCTTGCATTAA ATATCCCTCAA TAAACAGCCA	e289	e316	e61	e363
N189	AGGAGTAAACT TGTGCTTGA TCTTGTGTGT AGTTGGTAGTA	e364	e318	e289	e365
B190	AACCTCGAAA TCCTAGACATT AGTTTACTCCT ATCACCAACAA	e131	e320	e364	e59
H191	TTTTTTTTTT AGCTGATTGA TTTCGAGGTT AGAACAGGAT	nullT11S	e321	e131	e368
N192	ACTGAGGATAAA ATAACGGTGA TTTTTTTTT AAAAGTGAACA	e250	e323	nullT10N	e370
N193	AGCCATCTTT TTCCCTCACTAT TTATCCTCAGT AGCCTTCTT	e371	e325	e250	e27
N194	ACAGTAGGATT TTGGTGGTAA AAAGATGGCT TGTAGGAATCT	e157	e327	e371	e374
S195	AACACCTCAT ACCAAATGGAAA AATCCTACTGT TGCCATATCA	e375	e329	e157	e376
H196	AGAACTAAGGA AAGGGACAAT ATGAGGTGTT TATGACTGTGA	e136	e331	e375	e378
S197	AACTTCTCGA ACCTCTAAAGT TCCTTAGTTCT AAGACCAACA	e3	e180	e136	e380
H198	AGGAGTTACTT AGTTGATGCT TCGAGAACGTT TGAAAGAGGT	e381	e335	e3	e357
S199	TCGCTTTCTA TACCTTCTGA AAGTAACCTCT TCCAATCCAA	e383	e42	e381	e143
H200	TATGGAGTTCT AGCAGGATT TAGAAAGCGA AGAAGTAGACA	e385	e339	e383	e386
B201	TCCACGTTAT ATGTCACCT AGAACCTCCATA AACCGAAACT	e387	e341	e385	e388

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
H202	AGGGATTTAGT AGAGCGTAAT ATAACGTGGA AAAGCATTGAT	e389	e343	e387	e390
S203	ACGAACCATA TATCTTCCTGT ACTAAATCCCT AGCACTTACA	e391	e345	e389	e310
H204	TAACTGGAAGA AAATGAGGCT TATGGTCGT AAAATGGACA	e393	e347	e391	e394
N205	ACATCCATGA TCCTAGACATT TCTTCAGTTA ATTCACTCGA	e151	e320	e393	e396
N206	AATAGGGTACA TGAAAGTCGA TCATGGATGT TTCTGTAGTGA	e93	e351	e151	e398
N207	AAAAAACGCA TTTCTTTGCT TGTACCTATT TTCTTCACA	e399	e177	e93	e400
H208	AGTAATCTGGT AATGAGAGCT TGCCTTTTT ATTGTTGACA	e401	e273	e399	e38
N209	ATCACGACTT TACCTCTTCA ACCAGATTACT TTGCACCTTA	e403	e357	e401	e404
H210	TCGAGAGATTA TTAGCTTGCT AAGTCGTGAT TTTACTGGGTA	e405	e359	e403	e406
N211	AATCCGAACT AGCTACCATAT TAATCTCTCGA TTCTGCTCAA	e248	e89	e405	e408
H212	TTCTATGTGGA TGGCTGTTAA AGTTCGGATT TTATGTTGCA	e213	e363	e248	e410
S213	ATTGCCTACA TACTACCAACT TCCACATAGAA TCGTCTTCAT	e411	e365	e213	e412
H214	AGGAAGGAATA TTGTGGTGT TGTAGGCAAT TGATTAGGAGT	e270	e59	e411	e414
S215	TTTTTTTTT ATCCTGATTCT TATTCCCTCCT AACGTCTATCT	nullT10S	e368	e270	e415
S216	TTGTTCCCTT TGTTCACTTT TTTTTTTTTT ATCTTCCGAA	e416	e370	nullT11N	e417
H217	TAATGGGAGA AAGAAAGGCT AAGGGAACAA AGATCTGAGTT	e418	e27	e416	e419
N218	ATCTGCTTCA AGATTCTACA TCTCCCATTTA ACTTGCCTAA	e420	e374	e418	e237
H219	TTGGTTTGAAT TGATATGGCA TGAAGCAGAT ATAGCAGGATA	e422	e376	e420	e423
N220	AAACTCCGAA TCACAGTCATA ATTCAAACCAA ACATCGCTAA	e39	e378	e422	e425
H221	TTTGATTCTGT TGTTGGTCTT TTCGGAGTTT ATATAGAGGCA	e332	e380	e39	e427
S222	TTTCGTCTCA TACCTCTTCA ACGAATCAAAA TAAGCCACAA	e428	e357	e332	e429
H223	ACTATGTGAGT TTGGATTGGA TGAGACGAAA ATTACGAGAGA	e430	e143	e428	e431
S224	AGCCATCTT TGTCTACTTCT ACTCACATAGT TAACACAGCA	e371	e386	e430	e433
H225	TAGCATGAGTA AGTTCTGGTT AAAGATGGCT AATGTCTAGGA	e434	e388	e371	e320
S226	ACACGACTAA ATCAATGCTTT TACTCATGCTA ATGACCCAAT	e436	e390	e434	e224
H227	TAGTGGTTGTA TGTAAGTGTCT TTAGTCGTGT TTTTTAACGCT	e438	e310	e436	e70
S228	TTTACTGCCT TGTCCATTCTT TACAACCCTA ACATACACGT	e440	e394	e438	e441
N229	AAGTTAGGGTA TCGAGTGAAT AGGCAGTAAA AAAATGCAAGA	e84	e396	e440	e316
B230	TTCCAACAGA TCACTACAGAA TACCCCTAACCTT ACGTTAACCT	e301	e398	e84	e445
H231	TAAAATTGGCA TGTGAAGGAA TCTGTTGGAA TACATGGAAAGA	e319	e400	e301	e447
S232	AGCCTACAAA TGTCAAACAAAT TGCCAATTAA ACCAACAGA	e114	e38	e319	e449
N233	AGAGAATGAGT TAAGGTGCAA TTGTAGGCT TAGTTGTGAGA	e450	e404	e114	e451
S234	TTCCCAAAGA TACCCAGTAAA ACTCATTCTCTT AATCGTTCCCT	e187	e406	e450	e12
B235	AGTGTGTTGAA TTGAGCAGAA TCTTTGGAA TAAGAAGGTCA	e454	e408	e187	e231
N236	ATCGTAACCA TGCAAACATAA TTCAAAACACT ACATAACCGA	e268	e410	e454	e457
H237	TAGCTAATGGA ATGAAGACGA TGGTTACGAT TAGTTGTGAGA	e458	e412	e268	e451
N238	TACCGTCATT ACTCCTAATCA TCCATTAGCTA TCCAATCCAA	e460	e414	e458	e143
B239	TTTTTTTTT AGATGACGTT AATGACGTTA AGAGTTATGGAA	nullT11S	e415	e460	e462
H240	AGAGTGTGAT TTCGGAAGAT TTTTTTTTTT TTGTTGCATAA	e463	e417	nullT10N	e464
S241	ACTTGTTCCTT AACTCAGATCT ATCATCACTCT ATCGACCTTT	e54	e419	e463	e466
N242	TGAGTATCTGT TTAGGCAAGT AGGAACAAGT TGAAGTAAGGA	e467	e237	e54	e271
S243	ATATGCACCA TATCCTGCTAT ACAGATACTCA ATTCACTCGA	e223	e423	e467	e396
N244	AGGGTTCTAAT TTAGCGATGT TGGTGCATAT TAGGGTAACAT	e161	e425	e223	e117
S245	ACAACTCACT TGCCTCTATAT ATTAGAACCCCT ACATAACCGA	e473	e427	e161	e457
H246	AGTCGATGATA TTGTGGCTTA AGTGAGTTGT ATGAGTTAGGT	e475	e429	e473	e476
S247	ACAGCCATT TCTCTCGTAAT TATCATCGACT AATCTCCGAA	e46	e431	e475	e478
H248	ATGGATTAGGA TGCTGTGTTA TAATGGCTGT ACAATTGGAAA	e479	e433	e46	e480
S249	TCAGCATCTT TCCTAGACATT TCCTAATCCAT ACACCATCAT	e481	e320	e479	e204
B250	TTTGTTTCAGT ATTGGGTCTAT AAGATGCTGA TGGAGTAAGAT	e483	e224	e481	e186
N251	AATTCCCAGT ACGCTTAAAAAA ACTGAACAAAA ATAACGCTCT	e485	e70	e483	e486
H252	TTGGGATATCT ACGTGTATGT ACTGGGAATT TAATACGAGGA	e487	e441	e485	e74
N253	ATACTGCCAA TCTTGCATTAA AGATATCCCAA TCTTGTCA	e489	e316	e487	e490
H254	TGGATGATACT AGGTTAACGT TTGGCAGTAT TATAGACGTGT	e41	e445	e489	e255
N255	ACTTTCTCGT TCTTCCATGTA AGTATCATCCA ATCGCTACAA	e493	e447	e41	e494
H256	TGGATAGCTTA TCTGTTGGT ACGAGAAAGT ATATAGGACGT	e311	e449	e493	e496
N257	TTCATCCCCAA TCTCACAACCTA TAAGCTATCCA ATCCAACCAT	e497	e451	e311	e498
H258	AATAGGGTACA AGGAACGATT TTGGGATGAA TGGAATAGTCA	e93	e12	e497	e500
S259	ACAACTTCGA TGACCTTCTTA TGTACCTATT AGTCCAACAA	e326	e231	e93	e502
H260	AGTAGATACGT TCGGTTATGT TCGAAGTTGT AAGGTAGTCAT	e503	e457	e326	e504

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
S261	ACACGACTAA TCTCACAACTA ACGTATCTACT AAAGCACTCT	e436	e451	e503	e506
N262	TAGAGGAAAGT TTGGATTGGA TTAGTCGTGT TTTGGTCAA	e153	e143	e436	e113
S263	TTTTTTTTT TCCATAACTCT ACTTTCCCTCA ATTCAACCT	nullT10S	e462	e153	e509
S264	TTCACCACTT TTATGCAACAA TTTTTTTTT TCATAGCCAA	e510	e464	nullT11N	e167
H265	TGGCTTTTTTA AAAGGTCGAT AAGTGGTGAA ATTACGAGAGA	e512	e466	e510	e431
N266	AAACCCATGA TCCTTACTTCA TAAAAAAAGCCA AAACGTCTCA	e514	e271	e512	e515
B267	ATTAATTGCGA TCGAGTGAAT TCATGGGTTT AAAGAGAGGAA	e516	e396	e514	e222
S268	ATCGACATCA ATGTTACCTCA TCGCAATTAAAT TCTGACAACA	e518	e117	e516	e519
B269	AGGGTTAGAAT TCGGTTATGT TGATGTCGAT ACAGGAAGATA	e520	e457	e518	e345
S270	TGCTCCATTA ACCTAACTCAT ATTCTAACCCCT ATCACCCAAA	e178	e476	e520	e4
H271	TATGACAGAGT TTCGGAGATT TAATGGAGCA ATAGGACATGA	e524	e478	e178	e525
S272	TCGATCAACA TTTCAATTGT ACTCTGTCATA TATCGCTCAA	e526	e480	e524	e527
N273	AGACTAATGGA ATGATGCTGT TGTGATCGA TGGAGTAAGAT	e528	e204	e526	e186
N274	TTCACCACTT ATCTTACTCCA TCCATTAGTCT ACCATCCAAA	e510	e186	e528	e306
N275	AGTCTAAAGGA AGAGCGTTAT AAGTGGTGAA AGAAGTAGACA	e532	e486	e510	e386
S276	AACGCAATTTC TCCTCGTATTAA TCCTTAGACT ACTAGCACAA	e534	e74	e532	e19
H277	TGTTGAACTT TGTGACAAGA AAATTGGTTT AGGAGCATATA	e133	e490	e534	e537
S278	ACATTCACTT ACACGTCATAA AAGTTCAAACAA ACTAACGACA	e538	e255	e133	e156
N279	TTATCTGTGGA TTGTAGCGAT AGGTGAATGT TATTGAGGCTA	e125	e494	e538	e541
S280	ACCAACTTGA ACACGTCCTATAT TCCACAGATAA ATCCAACGTA	e542	e496	e125	e543
H281	ACAGGTTAGAT ATGGTTGGAT TCAAGTTGGT ACAGGAAGATA	e544	e498	e542	e345
N282	TACAACGTCA TGACTATTCCA ATCTAACCTGT AATCCGTTCT	e546	e500	e544	e547
N283	TAAGAGTGTGA TGTTGAACT TGACGTTGTA TTAAGGGAGAA	e548	e502	e546	e549
S284	TTTCGTCTCA ATGACTACCTT TCACACTCTTA ACGATCTCAA	e428	e504	e548	e551
H285	TAGGAAAGTCA AGAGTGCCTT TGAGACGAAA TTAGATACGGA	e552	e506	e428	e553
N286	TAACAACCGT TTGAACCAAAA TGACTTTCCCTA ATGACCCAAT	e554	e113	e552	e224
H287	TTTTTTTTT AGGTTGGAAT ACGGTTGTTA ATCAGGATGTA	nullT11S	e509	e554	e556
N288	TTCGAAATGTT TTGGCTATGA TTTTTTTTT TTGTTGCATAA	e557	e167	nullT10N	e464
S289	AACCTGAACA TCTCTCGTAA AACATTGCAA AAATCCGACT	e559	e431	e557	e560
H290	AGGTATCGTAT TGGACAGTTT TGTCAGGTT AGGTTAGACAT	e315	e515	e559	e341
S291	ACCTGACATT TTCCCTCTTTT ATACGATACCT ACATAACCGA	e256	e222	e315	e457
H292	ACGTTGATTTT TGTTGTCAGA AATGTCAGGT TAAAGAGGGAA	e565	e519	e256	e566
S293	AACTCGCTAA TATCTTCCTGT AAAATCAACGT TTCTCTGCT	e567	e345	e565	e568
H294	AACGTGAAATT TTTGGGTGAT TTAGCGAGTT TCTAAGTGTGA	e258	e4	e567	e173
B295	TCTTCACGT TCATGCTCTAT AATTTCACGTT ACTTGACACA	e43	e525	e258	e572
N296	ATTGCTGATT TTGAGGATA ACGTGAAGA TTTAAATGCGA	e573	e527	e43	e190
S297	TTCAGCTCAA ATCTTACTCCA AATCAGCAAAT ATTTCAGCCT	e575	e186	e573	e576
H298	AGTTGATAGGT TTTGGATGGT TTGAGCTGAA TGAGTAGAACAA	e189	e306	e575	e578
B299	ATCCACCATT TGTCTACTTCT ACCTATCAACT AACGTCTACT	e215	e386	e189	e415
B300	ACATATGGAGT TTGTGCTAGT AATGGTGGAT TTTCATTGGT	e581	e19	e215	e329
S301	ACCCCTCAAT TATATGCTCT ACTCCATATGT ATTCACTCGA	e71	e537	e581	e396
H302	TTTTGATTCTGT TGTCGTTAGT ATTGAAGGGT AGGATATAGCA	e332	e156	e71	e94
S303	AACTGCTCTT TAGCCTCAATA ACGAACCAAAA ACGATCTCAA	e587	e541	e332	e551
H304	AGTAGTGACAT TACGTTGGAT AAGAGCAGTT AAGAAAAGCAA	e589	e543	e587	e590
S305	TCCACTCAAA TATCTTCCTGT ATGTCACTACT AGCCTCATTT	e591	e345	e589	e347
H306	TGGCTTTTTTA AGAACGGATT TTTGAGTGGAA AGTGAAGTAAGA	e512	e547	e591	e594
S307	ACAGCCATTAA TTCTCCCTAA TAAAAAAAGCCA TCAACCTGTT	e46	e549	e512	e596
N308	TCGGAATTAGA TTGAGATCGT TAATGGCTGT AGATAGCGTAT	e597	e551	e46	e598
N309	AGACAACCTT TCCGTATCTAA TCTAATTCCGA AAACTGCTCA	e599	e553	e597	e515
H310	TAAGAGTGTGA ATTGGGTGAT AAGGTTGTT TAAATGTGCA	e548	e224	e599	e602
S311	TTTTTTTTT TACATCCTGAT TCACACTCTTA ACGATCAACT	nullT10S	e556	e548	e603
S312	AGCCTACAAA TTATGCAACAA TTTTTTTTTT TCAAGAACCA	e114	e464	nullT11N	e253
H313	AGACGGTAATA AGTCGGATT TTTGTAGGCT TTCTGTAGTGA	e606	e560	e114	e398
S314	TTACGTCAA ATGTCCTAACCT TATTACCGTCT ACGATCTCAA	e608	e341	e606	e551
H315	ATGAAAATGCT TCGGTTATGT TTGGACGTTAA TAAAGGCAAA	e610	e457	e608	e6
S316	AAACCTGCTA TCCCTCTTTAA AGCATTTCAT TCACAAACCT	e342	e566	e610	e613
H317	AAAAGGGTCTA AGCAGAAGAA TAGCAGGTTT AGTGGTAGTA	e614	e568	e342	e365
S318	TTAACGACCT TCACACTTAGA TAGACCTTTT TATCAACGCT	e616	e173	e614	e36
H319	TCGTAGTATGT TGTGTCAAGT AGGTGCTTAA AGAGAAGATGT	e618	e572	e616	e619

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
B320	AGACCACTTT TCGCATTAAA ACATACTACGA TACACAAGCT	e620	e190	e618	e621
H321	AGAGAAATGACT AGGCTGAAT AAAGTGGTCT ATTAGTCTGGA	e450	e576	e620	e623
S322	ACCAGCATAA TGTTCTACTCA ACTCATTCTCT TTCTGCTCAA	e624	e578	e450	e408
H323	TCGGAATTAGA AGATGACGTT TTATGCTGGT ATTATCGGAGA	e597	e415	e624	e627
S324	TCGCATTCTA ACCAAATGAAA TCTAATTCCGA AAGTCCCCAT	e628	e329	e597	e629
H325	TAGATGTACGT TCGAGTGAAT TAGAATCGA TTAGAGTGACA	e630	e396	e628	e631
S326	AATACCACGA TGCTATATCCT ACGTACATCTA AAACGTGCCA	e632	e94	e630	e515
H327	TGAGTATCTGT TTGAGATCGT TCGTGGTATT TAGGAGAAACT	e467	e551	e632	e635
S328	TACAACGTCA TTGCTTTCTT ACAGATACTCA ACCAACGCTAA	e546	e590	e467	e359
H329	AGAAGGATGAT AAATGAGGCT TGACGTTGTA AGGTTAGGTA	e638	e347	e546	e639
N330	AAACCTGCTA TCTTACTCACT ATCATCCTCTT TTACACCTGTT	e342	e594	e638	e641
H331	AGAGATGACTT AACAGGTTGA TAGCAGGTTT AGAGTTGAGTA	e275	e596	e342	e643
S332	AACCCCTTTT ATACGCTATCT AAGTCATCTCT AAGACCACAT	e155	e598	e275	e645
H333	TTTAGTGGAGA TGGACAGTTT AAAGAGGGTT ACTAGATGGAA	e646	e515	e155	e647
S334	ACGACCAATA TGCACATTTA TCTCCACTAAA AACGTACCTT	e648	e602	e646	e649
H335	TTTTTTTTTT AGTTGATCGT TATTGGTCGT ATAGGACAAGT	nullT11S	e603	e648	e650
H336	TTCGGATAGAA TGGGTTCTGA TTTTTTTTTT TGGGTTATAT	e651	e253	nullT10N	e652
S337	ACCTTCAGTT TCACTACAGAA TTCTATCGAA AAGTCCCCAT	e653	e398	e651	e629
B338	TGAGTAGGATT TTGAGATCGT AACTGAAGGT AAATTGGAACA	e655	e551	e653	e102
N339	TCGATCAACA TTTGCCTTTA AATCCTCTACA ACACCTCCAT	e526	e6	e655	e658
H340	AGGACTAATGT AGGTTTGTGA TGTTGATCGA TAAAACGTGTT	e659	e613	e526	e660
S341	ACCAACAAGA TACTACCAACT ACATTAGTCCT AACACCTCCTT	e661	e365	e659	e179
H342	TGGAAGAGTAT AGCGTGTGATA TCTTGTGGT AGGTTCTAGTT	e101	e36	e661	e145
S343	AGCACCTTTA ACATCTTCTCT ATACTCTTCCA TTCCAGTTCA	e665	e619	e101	e666
H344	TTTGTTCAGT AGCTTGTGTA TAAAGGTGCT ATAGTGAGGAA	e483	e621	e665	e325
S345	TACACGAACA TCCAGACTAAT ACTGAACAAAA TCCCAGAAAA	e669	e623	e483	e670
H346	TTCGAAATGTT TTGAGCAGAA TGTCGTGTA TGAAGTAAGGA	e557	e408	e669	e271
N347	ACATTCACCT TCTCCGATAAT AACATTGCGAA ATAACGCTCT	e538	e627	e557	e486
H348	AGACTAATGGA ATGGGAACCTT AGGTGAATGT ATTGTAGGACT	e528	e629	e538	e676
S349	AACGCAATTGTT GTGCACTCTAA TCCATTAGTCT TCATGCCCTA	e534	e631	e528	e294
N350	TGCAGTTTTA TGGACAGTTT AAATTGGCGTT AGATCTGAGTT	e48	e515	e534	e419
N351	TTTACCGACA AGTTTCTCCTA TAAAAACTGCA TCCCAATTGA	e681	e635	e48	e682
H352	TAGGAACAGAA TTAGCTTGGT TGTCGGTAAA TTTAAATGCGA	e344	e359	e681	e190
S353	ACCAACATAA TACCTAAACCT TTCTGTTCCCTA ATTCTGCACT	e624	e639	e344	e686
H354	ACAAGGATAGA AACAGGTGAA TTATGCTGGT TGGGTACTTTA	e116	e641	e624	e688
S355	AGACCACTTT TACTCAACTCT TCTATCCTGT ATCGCTACAA	e620	e643	e116	e494
H356	TAAAATTGGCA ATGTTGGCTT AAAGTGGCT TAAGAAGGTCA	e319	e645	e620	e231
N357	AGCCATCTTT TTCCATCTAGT TGCCAATTTTA ACAAGTCCAT	e371	e647	e319	e694
H358	AGTTGTAGTCT AAGGTACGTT AAAGATGGCT TACATGATGGA	e695	e649	e371	e10
S359	TTTTTTTTT ACTTGTCTTAT AGACTACAAC AGAAACCCAA	nullT10S	e650	e695	e697
B360	TCAATCTCGT ATATACACCCA TTTTTTTTTT TCAAGACACA	e698	e652	nullT11N	e318
H361	ATTGCAATGAT ATGGGAACCTT ACGAGATTGA TAGGTTAACAT	e700	e629	e698	e117
B362	TCCATATCCCT TGTTCCAATT ATCATTGCAT ACACACTCAA	e702	e102	e700	e703
H363	AGGTAATGGTA ATGGAAGTGT AGGGATAGGA TGCAATAGGT	e704	e658	e702	e705
S364	AACTTCAGCT AACACGTTTA TACCATTAACCT TAACTGCCAA	e99	e660	e704	e707
N365	AACTGTAGGA AAGGAGGTTT AGCTGAAGTT TGAGGTATGAT	e708	e179	e99	e45
S366	TTCGTACACA AACTAGAACCT TCCTATCAGTT AGCCTCATTT	e63	e145	e708	e347
H367	TTAAGGACTGA TGAACGGAA TGTGTACGAA AGAATATGGGT	e712	e666	e63	e713
S368	ATTCGCTACT TTCCTCACTAT TCAGTCCTAA TTTTTTTTTT	e714	e325	e712	nullT10E
H369	TGAGGATGTTA TTTTCTGGGA AGTAGCGAAT TTTTTTTTTT	e715	e670	e714	nullT11E
S370	TCACACAGTT TCCTTACTTCA TAACATCCTCA TTTTTTTTTT	e716	e271	e715	nullT10E
H371	AGTTCTGAGAT AGAGCGTTAT AACTGTGTGA TTTTTTTTTT	e717	e486	e716	nullT11E
N372	ACACATCCAT AGTCCTACAAT ATCTCAGAACT TTTTTTTTTT	e718	e676	e717	nullT10E
H373	AAAGAGGAGAT TAAGGCATGA ATGGATGTGT TTTTTTTTTT	e719	e294	e718	nullT11E
S374	TCCATCGAAA AACTCAGATCT ATCTCCTCTTT TTTTTTTTTT	e720	e419	e719	nullT10E
H375	TGAGTATGAGT TCAATTGGGA TTTCGATGGA TTTTTTTTTT	e266	e682	e720	nullT11E
S376	TACCGTCATT TCGCATTAAA ACTCATCTACA TCAAGACACA	e460	e190	e266	e318
H377	AGAAAGTCAGAT AGTGCAGAAAT AATGACGGTA TAAGTGAACGTA	e352	e686	e460	e725
N378	ACCCCTTGAT TAAAGTACCCA ATCTGACTTCT ACATCGCTAA	e726	e688	e352	e425

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
N379	TGGCTATTAGA TTGTAGCGAT ATCAAAGGGT AGAAAGGAAGA	e65	e494	e726	e729
B380	ATCGCAACTA TGACCTTCCTA TCTAACATGCCA ATCGTCCAAT	e730	e231	e65	e731
H381	AGTGTGTTGAA ATGGACTTGT TAGTTCGAT TGAGAGAAAGA	e454	e694	e730	e733
S382	TTCTCTTGCT TCCATCATGTA TTCAAAACACT AACACTCGAT	e734	e10	e454	e735
B383	TTTTTTTTTT TTGGGTTCT AGCAAGAGAA AGTCATTAGT	nullT11S	e697	e734	e736
H384	AGTCTAAAGGT TGTGTCTTGA TTTTTTTTT AGTGTAAAGAGA	e737	e318	nullT10N	e738
S385	TTCACGAACT ATGTTACCTA ACCTTAGACT ACTCAACGTA	e739	e117	e737	e740
H386	TCGGTAGAGTA TTGAGTGTG AGTCGTGAA ATGGTAGATCA	e741	e703	e739	e742
B387	ATCTGCTTCA ACCTATATGCA TAATCTACCGA ACACCATCAT	e420	e705	e741	e204
H388	ACATATGGAGT TTGGCAGTTA TGAAGCAGAT TTAGAGTGTCT	e581	e707	e420	e746
S389	AAACCCAAGA ATCATACCTA ACTCCATATGT AAAACGCAAT	e747	e45	e581	e44
B390	TGGAAGAGTAAATGAGGCT TCTTGGGTTT AATGTGGAAAA	e101	e347	e747	e141
S391	TTTTTTTTT ACCCATATTCT ATACTCTTCCA AGAACCCAAA	nullT10S	e713	e101	e751
B392	TATATGGTCGT TGTGTCTTGA TTTTTTTTT TATCGGAGAAA	e193	e318	nullT10N	e753
S393	AAACCCATGA TCAGTCACTTA ACGACCATA ATA CGCTACAA	e514	e725	e193	e494
H394	TATGGAGCTAT TTAGCGATGT TCATGGGTTT AGGAAATAGGA	e756	e425	e514	e757
S395	TGATCCACAT TCTTCCTTCT ATAGCTCCATA ATGCATACCA	e758	e729	e756	e124
H396	TGTAGAGGATT ATTGGACGAT ATGTGGATCA AGGAGCATATA	e655	e731	e758	e537
S397	AACGTCTTCT TCTTTCTCTA AATCCTCTACA TTCGATCACT	e142	e733	e655	e763
H398	TTTTGTCTGT ATCGAGTGTG AGAAGACGTT ATAGGACATGA	e764	e735	e142	e525
S399	TTTTTTTTT ACTAATGACCT ACAGACAAAAA ATCTTCTGCA	nullT10S	e736	e764	e135
B400	AGCTTCATCA TCTCTTACACT TTTTTTTTTT TTCCCTTCCA	e146	e738	nullT11N	e768
H401	TAGAAGGCATA TACGTTGAGT TGATGAAGCT TATATGACGGA	e307	e740	e146	e210
S402	ATTCCGTACA TGATCTACCAT TATGCCCTCTA TTCTGACCAA	e771	e742	e307	e184
B403	AGAATTAGGGA ATGATGGTGT TGTACGGAAT TAGGGTAACAT	e108	e204	e771	e117
S404	AACTTCTCGA AGACACTCTAA TCCCCTAATTCT AACACAGACA	e3	e746	e108	e776
H405	AGTGGATCTTA ATTGCGTTT TCGAGAAAGTT TGTGACAGATA	e777	e44	e3	e778
S406	TAACAACCGT TTTTCCACATT TAAGATCCACT ACATTCCACT	e554	e141	e777	e780
H407	TTTTTTTTT TTTGGGTTCT ACGGTTGTTA TTGACGAAAAT	nullT11S	e751	e554	e781
S408	ACCATGTCAT TTTCTCCGATA TTTTTTTTTT TCAAGAACCA	e782	e753	nullT11N	e253
H409	AGGTGAAGTAT TTGTAGCGAT ATGACATGGT ATGACAGTAGT	e784	e494	e782	e785
S410	TTCGCTCTAA TCCTATTCTCT ATACTTCACCT TCACAACGTA	e309	e757	e784	e282
B411	TGAGGACTATT TGGTATGCA TTAGAGCGAA TAGAGGAACAT	e76	e124	e309	e789
S412	TTTACTGCCT TATATGCTCT AATAGTCTCTA TCATGACACA	e440	e537	e76	e791
H413	AGGGATTCTATA AGTGTGCGAA AGGCAGTAAA TACATGATGGA	e792	e763	e440	e10
S414	TTCTCTTGCT TCATGCTCTAT TATGAATCCCT AGTCCAACA	e734	e525	e792	e502
H415	TTTTTTTTT TGCAGAAGAT AGCAAGAGAA ATTACAGAGGT	nullT11S	e135	e734	e796
H416	AGGGATGTAAT TGGAAGAAA TTTTTTTTTT AGAGTGGATAA	e88	e768	nullT10N	e798
S417	TGCTACACAA TCCGTCTATATA ATTACATCCCT ACACCATCAT	e799	e210	e88	e204
H418	AAAGTGAGGATA TTGGTCAGAA TTGTGTAGCA AAAAATGGACA	e801	e184	e799	e394
B419	ATTCCTGCTACA ATGTTACCTA TATCCTCACTT AATCCGTTCT	e771	e117	e801	e547
H420	ATTTGCTGATT TGTGTGTGTT TGTACGGAAT TTAAGGAAGGT	e573	e776	e771	e806
S421	AAACCCAAGA TATCTGTCACA AATCAGCAAAT ACAAAAGTCCA	e747	e778	e573	e808
H422	AGTATGTCGTGA AGTGGAAATGT TCTTGGGTTT TCAGTTAGGA	e809	e780	e747	e810
B423	TTTTTTTTT ATTTTCGTCAA TCAGACATACT AAAGACACCT	nullT10S	e781	e809	e811
N424	AGTAGATACGT TGGTCTTGA TTTTTTTTT TATCGGAGAAA	e503	e253	nullT10N	e753
S425	ACCACTGATT ACTACTGTCAT ACGTATCTACT AAACGCAAAT	e814	e785	e503	e815
H426	TGGGATCTATT TACGTTGTGA AATCAGTGGT TATACGAAGGT	e816	e282	e814	e817
B427	TACACGTCTAT ATGTTCTCTA AATAGATCCCA ATCGACTCAA	e818	e789	e816	e819
H428	AGGTAAGAGAA TGTGTCTGTA ATGACGTGTA AGGCTTGATA	e820	e791	e818	e821
N429	AGCTCAATCA TCCATCATGTA TTCTCTTACCT ACGTTAACCT	e822	e10	e820	e445
H430	TATGGAGCTAT TGTGGAACAT TGATTGAGCT AGAAACTATGGT	e756	e502	e822	e154
S431	TTTTTTTTT ACCTCTGTAAT ATAGCTCCATA TCAAAACGACT	nullT10S	e796	e756	e826
S432	TATTCTCGCA TTATCCACTCT TTTTTTTTTT ACAGCATACT	e827	e798	nullT11N	e828
H433	AGGATGTACTT ATGATGGTGT TGCAGAGATA ACAATTGGAAA	e829	e204	e827	e480
S434	ACCACTGTTA TGTCCATTAA AAGTACATCCCT TCGACTTTCA	e831	e394	e829	e351
H435	AACAGAGAGAT AGAACGGATT TAAAGCTGGT TATTGAGGTA	e833	e547	e831	e834
S436	ATTCGACCAT ACCTTCCTAA ATCTCTGTGTT AGCCATACAT	e835	e806	e833	e836
H437	AGGTATGGAAT TGGACTTTGT ATGGTGAAT TGGAGTAAGAT	e837	e808	e835	e186

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
S438	ACGATCATCA TCCTAAACTGA ATTCCATACCT ATTCTCAGCT	e313	e810	e837	e840
H439	TTTTTTTTTT AGGTGTCCTT TGATGATCGT ATTTGCAGAT	nullT11S	e811	e313	e841
S440	TTCCCCAAGA TTTCTCCGATA TTTTTTTTTT ACGTTCACTA	e187	e753	nullT11N	e843
H441	TTTGACTGTTT ATTTGCGTT TCTTTGGAA TTTTCGATTGT	e844	e815	e187	e845
S442	TTCTACGCA ACCTTCGTATA AAACAGCTAAA TACAAGCCAA	e58	e817	e844	e847
H443	TGAGTATCTGT TTGAGTCGAT ATGCGTAGAA TAGAGGAACAT	e467	e819	e58	e789
S444	TCGCATCTA TATCAAGACCT ACAGATACTCA ACTAGCACAA	e163	e821	e467	e19
H445	AGATGGAGATT AGGTTAACGT TAGATCGAA TAGGTTTCAGA	e852	e445	e163	e853
S446	TAACTCGCTT ACCATAGTTCT AATCTCCATCT ATCCAACGTA	e854	e154	e852	e543
H447	TTTTTTTTTT AGTCGTTGA AAGCGAGTTA AGGTATTAGT	nullT11S	e826	e854	e736
H448	ATCTTAGGGAA AGTATGCTGT TTTTTTTTTT AGAGTAAGGTT	e857	e828	nullT10N	e858
S449	TTTCCAGTC A TTTCAATTGT T T C C T T A A G A T T C C A T C A C T	e859	e480	e857	e763
H450	AGGAGTGTATT TGAAAGTCGA TGACTGGAAA ACAGGAAGATA	e861	e351	e859	e345
S451	ATTTCCTCGT TACCTCGAATA AATACACTCCT TCCACAAGTT	e863	e834	e861	e864
H452	TTGGGATATCT ATGTATCGCT ACGGAGAAAT ACTTTAGAGGT	e487	e836	e863	e180
S453	ATCGCTAACAA ATCTTAACCTCA AGATATCCCAA TCACAACGTA	e867	e186	e487	e282
H454	TGGCTATTAGA AGCTGAGAAT TGTTAGCGAT TGTAGGAATCT	e65	e840	e867	e374
S455	TTTTTTTTTT ATCTGCAAAAT TCTAATAGCCA ATTCAACCGTA	nullT10S	e841	e65	e90
H456	TTGGTTTGAAT TAGTGAACGT TTTTTTTTTT AGGCTAAGTAT	e422	e843	nullT10N	e873
S457	ACCTCACATT ACAATCGAAAA ATTCAAAACCAA ACCAAGCTAA	e67	e845	e422	e359
B458	TTATCTGTGGA TTGGCTTGT AATGTGAGGT ACGGAAATTTT	e125	e847	e67	e66
S459	ACGACCAATA ATGTTCCCTA TCCACAGATAA AGCATCAACT	e648	e789	e125	e335
B460	AGTCGATGATA TTGTGCTAGT TATTGGTCGT TCTGGGATTAT	e475	e19	e648	e235
S461	ACACACAAAGA TCTGAAACCTA TATCATCGACT AACCGAAAT	e289	e853	e475	e815
H462	TGATAGGACTT TACGTTGGAT TCTTGTGTGT TCAGTTAGGA	e884	e543	e289	e810
B463	TTTTTTTTTT ACTAATGACCT AAGTCCTATCA TATGCACTCA	nullT10S	e736	e884	e75
S464	ACAATCCGAT AACCTTACTCT TTTTTTTTTT TTTTTTTTTT	e887	e858	nullT11N	nullT10E
N465	TTAAGGACTGA AGTATCGAA ATCGGATTGT TTTTTTTTTT	e712	e763	e887	nullT11E
S466	ATCGACATCA TATCTTCCTGT TCAGTCCTAA TTTTTTTTTT	e518	e345	e712	nullT10E
H467	ATGGATCTGAT AACTTGTTGA TGATGTCAT TTTTTTTTTT	e890	e864	e518	nullT11E
N468	TACCAAAGCA ACCTCTAAAGT ATCAGATCCAT TTTTTTTTTT	e891	e180	e890	nullT10E
H469	AGTAGACTGAA TACGTTGTGA TGCTTTGGTA TTTTTTTTTT	e73	e282	e891	nullT11E
S470	TTACACAGAA AGATTCCCTACA TTCAGTCTACT TTTTTTTTTT	e893	e374	e73	nullT10E
B471	TTTTTTTTTT TACGGTGAAT TTCGTGGAA TTTTTTTTTT	nullT11S	e90	e893	nullT11E
S472	TCACACAGTT ATACTTAGCCT TTTTTTTTTT TTTTTTTTTT	e716	e873	nullT11N	nullT10E
N473	TTAGGATGACA TTAGCTTGGT AACTGTGTGA TTTTTTTTTT	e185	e359	e716	nullT11E
S474	ACAATCCGAT AAAATTCCGT TGTCACTCTAA TTTTTTTTTT	e887	e66	e185	nullT10E
H475	ATTCGTAGAGA AGTTGATGCT ATCGGATTGT TTTTTTTTTT	e897	e335	e887	nullT11E
S476	ATCTCATGCA ATAATCCCGA TCTCTACGAAT TTTTTTTTTT	e898	e235	e897	nullT10E
N477	ATGGGATCTGAT ATTTGCGTT TGCATGAGAT TTTTTTTTTT	e890	e815	e898	nullT11E
N478	ATCTCATGCA TCCTAAACTGA ATCAGATCCAT TTTTTTTTTT	e898	e810	e890	nullT10E
H479	TTTTTTTTTT TGAGTGCATA TGCATGAGAT TTTTTTTTTT	nullT11S	e75	e898	nullT11E
A480	ACAAGGGATAT TTTTTTTTTT TTTTTTTTT AGAGCTATGAT	e901	nullT10W	nullT10N	e182
A481	TAGCATCACA TTTTTTTTTT ATATCCCTGT TTGACACTT	e18	nullT11W	e901	e192
A482	TTAGTATCGGT TTTTTTTTTT AGGTAAAGCTT AGGAGCATATA	e9	nullT10W	e26	e537
A483	AGATGTCAGAT TTTTTTTTTT AAGAACAGT TAGGAGAAACT	e913	nullT10W	e11	e635
I484	TCACATACGT TTTTTTTTTT ATCTGACATCT TTCCATTGCA	e7	nullT11W	e913	e916
A485	TTAGGGAATCT TTTTTTTTTT ACGTATGTGA ATTAGTCTGGA	e917	nullT10W	e7	e623
I486	TTTCCCTTGA TTTTTTTTTT AGATTCCCTAA TAACACAGCA	e207	nullT11W	e917	e433
A487	AGGTTAACAGA TTTTTTTTTT TCAAGGGAAA TATTGAGGCTA	e5	nullT10W	e207	e541
I488	AAGATACGAGA TTTTTTTTTT ACGTATGTGA AGAGTTAGGAA	e20	nullT10W	e7	e194
A489	TTAACAGCTAGGT TTTTTTTTTT TAGAGCGAAA TCTAAGTGTGA	e929	nullT10W	e22	e173
A490	TTTCCCTTGA TTTTTTTTTT ACCTAGCTAA TACCATGCTT	e207	nullT11W	e929	e932
I491	TTTCGGATAGA TTTTTTTTTT TCAAGGGAAA AGTGGTAGTA	e28	nullT10W	e207	e365
I492	AAGATACGAGA TTTTTTTTTT TCAAGGGAAA TTTTTAACGCT	e20	nullT10W	e207	e70
I493	AACAAGTCCT TTTTTTTTTT TCTCGTATCTT TACAAGCCAA	e939	nullT11W	e20	e847
A494	TACGGTATTGA TTTTTTTTTT AGGACTTGT ACAGGAAGATA	e16	nullT10W	e939	e345
A495	TTTCGGATAGA TTTTTTTTTT TGTGATGCTA TGGAATAGTCA	e28	nullT10W	e18	e500
I496	AGGGATTCTATA AAGTGTGCAA AAGGTTCAAGT AAAAATGGACA	e792	e192	e110	e394

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
I497	TTTACTGCCT TCCCTATGAAA TATGAATCCCT TCATCACCAT	e440	e21	e792	e298
I498	ACATATGGACT AAGAAAGGCT AGGCAGTAAA AGGAGCATATA	e581	e27	e440	e537
A499	TGCAGTTTTA AGGAACGATT ATTGAAGGGT ATAGACAGGAA	e48	e12	e71	e57
A500	TTTGTTCACT TCGAATGAA TGTCGGTAA TTTGTCATGT	e483	e916	e681	e963
I501	TTTGATTCTG TGCTGTGTTA TGTCGCTGA ATTACGAGAGA	e332	e433	e669	e431
A502	TTGAGAGTAGT TGGATCGATT AAGAGCAGTT TTTCATTTGGT	e238	e8	e587	e329
A503	TCGTTTAGAGA TGGGTTCTT AAGGAATCGA TGTAGGAATCT	e217	e23	e240	e374
A504	TTCTATGTGGA AAGCATGGTA AAAGAGGGTT AGAAAGGAAGA	e213	e932	e155	e729
A505	TTTGTTCACT TAGGCAATGA TGTAGGCAAT AGTAATCAGGA	e483	e196	e411	e983
A506	AGTAGTGACAT TTGGCTTGT ACTGGGATT TTAAGGAAGGT	e589	e847	e485	e806
A507	ACAGGTTAGAT TTGTGCTAGT TTTGAGTGGA AGATAGCGTAT	e544	e19	e591	e598
A508	TTTTTTTTTG TGTGTTGAGT TGACGTTGTA TAATACGAGGA	nullT11S	e30	e546	e74
A509	AGGATGTACTT AGGAGGTAGA TTTTTTTTT AGTAGGAAGAA	e829	e229	nullT10N	e122
A510	TGAGGACTATT ATGGTGATGA TAAAGCTGGT ACGGAAATTTT	e76	e298	e831	e66
A511	AGCAGGTATTA TCAATTGGA AAAGAAGGGT AATGTGGAAAA	e1007	e682	e86	e141
I512	ACCAGCATAA ACATTGACAAA TAATACCTGCT TCGACTTCTT	e624	e963	e1007	e139
I513	AGTCGATGATA TTTTCTGGGA TTATGCTGGT TAATACGAGGA	e475	e670	e624	e74
A514	TAAATGGGAGA ATGTAGCACT TAGAATCGA TATAGACGTTG	e418	e241	e628	e255
I515	TATGGAGCTAT ATGTGACAGT TGAAGCAGAT TATTCGAGGTA	e756	e220	e420	e834
A516	TTATTTTGCCT ATGAAGACGA ATGTGGATCA TGAAAGAGGTA	e1027	e412	e758	e357
I517	ACCACTGATT TCCTGATTACT ACGCAAATAA TTAACAGCCA	e814	e983	e1027	e363
A518	AACAGAGAGAT AGAGCGTTAT AATCAGTGGT AAAGAGAGGAA	e833	e486	e814	e222
A519	AGAGATGACTT AAATGAGGCT ATGGTCAAT AATCGGAAAAAA	e275	e347	e835	e1036
I520	ATTAATTGCGA AGAACGGATT AAAGAGGGTT TACATGATGGA	e516	e547	e155	e10
I521	TTTTTTTTT TCCTCGTATTA TCGCAATTAAT ATTCCAACCT	nullT10S	e74	e516	e509
A522	TTGAGGGATAT TGTGTCATGA AAGGGTCAAT AGTACGAGTAT	e61	e791	e95	e1049
I523	TGGAAGAGTAT AAGAAAGTCGA TGATGAAGCT ATTGTTGACA	e101	e139	e146	e38
A524	TAGAGATGCTT TTCGGAGATT AGACTGGTTT TAAAACGTGTT	e1060	e478	e103	e660
I525	TCGATTCTT ATACGCTATCT AAGCCTCTCA ACTGTCACAT	e240	e598	e1060	e220
I526	TGTTGAACTT ATGGGAACTT AAGGAATCGA TATATGACGGA	e133	e629	e240	e210
A527	AGGAGTGTATT TTAGGCAAGT AGGTGAATGT ATTACGAGAGA	e861	e237	e538	e431
A528	TTTGATTCTG TGGTATGCAT ACGGAGAAAT AGTTGGTAGTA	e332	e124	e863	e365
A529	AGGAAGATTCT TGGCTGTTAA TGAGACGAAA AAATTGGAACA	e1076	e363	e428	e102
I530	ACTTCTCGT TTCCTCTCTT AGAATCTCCT TCCCAGAAAA	e493	e222	e1076	e670
A531	ACATATGGAGT ATGTATGGCT ACGAGAAAGT TGGTGGAGATTA	e581	e836	e493	e14
I532	ATTCATCGT TTTTCCGATT ACTCCATATGT TATCGCTCAA	e1082	e1036	e581	e527
A533	AGTGTGTTGAA ATGTGGCTT ACGATGGAAT TGTGACAGATA	e454	e645	e1082	e778
A534	TTTTTTTTT AGGTTGGAAT AGCAAGAGAA ATTGAAGCAAA	nullT11S	e509	e734	e274
A535	TTCGGATAGAA AAGGGTGAAT TTTTTTTT AGGAGTTACT	e651	e152	nullT10N	e17
A536	AGGTACGATTA TGGCTGTAAT AACTGAAGGT AGAGAAGATGT	e1093	e96	e653	e619
I537	AGCTTCATCA ATACTCGTACT TAATCGTACCT AGCTCTCATT	e146	e1049	e1093	e273
I538	AGAAACTAAGGA TGGCTGTTAA TGATGAAGCT AGCAAAGAAA	e136	e363	e146	e177
A539	TGGCTATTAGA TGGTTGCTA TTGTTGGCTA TTTCGATTGT	e65	e171	e138	e845
A540	AGGTAATGGTA TGGCAGTAAT AATGTGAGGT ACAGGAAGATA	e704	e104	e67	e345
A541	ATCTTGGAGTA ATGTGACAGT AGCTGAAGTT AGGAAATAGGA	e112	e220	e99	e757
A542	TTCGAAATGTT TGTCGTTAGT AATGTCAGGT AATGTGGAAAA	e557	e156	e256	e141
A543	AGGACTAATGT AACTGTGGA TGTCAGGTT TCTTAGAAGGT	e659	e864	e559	e218
A544	AATGCAGATTT TTGTGGCTA TCTTGTGGT ATATGGTAGCT	e328	e429	e661	e89
A545	TGGATGATACT TTTTCTGGGA TTTGGAGTGT TTAGATACGGA	e41	e670	e330	e553
A546	TAAAATTGGCA ATCGAGTGT TCTTGGGTTT AGGTTAGACAT	e319	e735	e747	e341
A547	TGGAAGAGTAT ATGGGAACTT TAATGGCTGT TTAGTAGGTGT	e101	e629	e46	e29
A548	AATAGGGTACA AATGAGAGCT TAAAGGTGCT AATTTTCGGT	e93	e273	e665	e202
A549	TTGGTTTGAAT AAGAAAGTCGA TCGCTTTTT ATGTAGTAGCA	e422	e139	e399	e259
A550	AGGGATTTAGT TGAAATGGGT AATGTGAGGT TGAAGTAAGGA	e389	e68	e67	e271
A551	AGGTGAAGTAT TTGGCACTTA TATGTTCTG AAGGTAGTCAT	e784	e707	e391	e504
A552	AGTGGATCTTA AGGTGTTCT TTAGAGCGAA TTAGTTCAGGA	e777	e257	e309	e263
A553	TAGTCAGTTGA AGTCGGATT ACGGTTGTTA TGAGAGAAAGA	e262	e560	e554	e733
A554	ACAAGGATAGA AAGGGAGGTTT AAGGTTCACT TGGAGACTAAT	e116	e179	e110	e296
A555	TCGGAATTAGA AAGGGACAAT TTGGTGACTT TAATACGAGGA	e597	e331	e118	e74

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
A556	AGAACAGAGTA ATTGCGTTT AAGGTTGTCT AGTATTACGGA	e234	e44	e599	e53
A557	AGACGGTAATA TGGACTTGT TGGATGTCAA TAGTTGTGAGA	e606	e808	e236	e451
A558	TTTTTTTTTT AGCTGATTGA TTGGACGTA ATTACAGAGGT	nullT11S	e321	e608	e796
A559	AGTAGTGAAGT ACGAGTTGAA TTTTTTTTT ACTTGGTAGAT	e56	e47	nullT10N	e1184
A560	TGTTGTTGATT TGAACGTGAA ATGCCTAGAA TAGGTTAACAT	e246	e666	e58	e117
A561	AGGGTTAGAAT TGTGAAGGAA AGTCGGATT AAAATGGACA	e520	e400	e248	e394
I562	TGATCCACAT TGCTACTACAT ATTCTAACCCCT AGTTCACACA	e758	e259	e520	e290
I563	AGGTATCGTAT TTAGCTTGGT ATGTGGATCA ACAGGAAGATA	e315	e359	e758	e345
A564	TAAGACTGTGA TGTAAGTGTCT TGATGATCGT TCTGGGATTAT	e548	e310	e313	e235
I565	TAGAAGGCCATA TACGTTGTGA TGAGACGAAA ACAGGAAGATA	e307	e282	e428	e345
A566	TGTAGAGGATT AGTGAATGT TTAGAGCGAA ACGGAAATTTT	e655	e780	e309	e66
A567	AGTCTAAAGGA ACGTGAAGAA TAGCAGGTTT ATTAGTGGGTA	e532	e119	e342	e49
I568	TGGATAGCTTA TTAGGCAAGT ATGGTGTCAA AATCGGAAAAAA	e311	e237	e82	e1036
A569	TATGGAGCTAT TTGAGATCGT TTGGATGAA AAAAGTTGTGT	e756	e551	e497	e206
I570	AAGCTCTTCA ATCTACCAAGT TTTTTTTTTT TCACCGTTAT	e1230	e1184	nullT11N	e323
A571	AGTCTAAAGGT TTGTGGTGAT TGAAGAGCTT TTTTCATGAT	e737	e59	e1230	e1233
A572	AAGAGGTAGAA ATGATGAGCA AGTCGTGAA ATAGTCTTGGA	e1236	e64	e739	e1237
I573	AACCATGTCT TGTCCATTTT TTCTACCTCTT ACCCAATGAT	e1238	e394	e1236	e1239
A574	ACGTTGATTT TGTGTGAAC AGACATGGTT TGGAGTAAGAT	e565	e290	e1238	e186
A575	ATTCTGTAGAGA TGTGTCTTGA TTAGCGAGTT ATAGGACAAGT	e897	e318	e567	e650
A576	TTAAGGACTGA TTGAGATCGT TGCTAGAGAT TTTTTTTTTT	e712	e551	e898	nullT11E
I577	TTAGGATGACA TGTAAGTGTCT TGATGTCAT TTTTTTTTTT	e185	e310	e518	nullT11E
A578	ATGGTTTTGTT AGAGCGTAAT AGTAGCGAAT TTTTTTTTTT	e356	e343	e714	nullT11E
A579	AGAGTGTGATTT TTGTGCTAGT TGTGATACGT ATTGATGCAA	e463	e19	e358	e1257
A580	AGAAGTACAGA TGTGGGTAAG AGGAACAAAGT AGGAAAGACTA	e1260	e83	e54	e280
I581	ACCCTTGAA TTTTCCGATT TCTGTACTTCT ACGATCTCAA	e1262	e1036	e1260	e551
A582	ACTGAGGATAA ATGGTTGGAT TTCAAAAGGGT AAAGGCTAGTA	e250	e498	e1262	e288
A583	TTTTTTTTT AGTCGTTGA TCTTGGGAA AGACTTATGGA	nullT11S	e826	e187	e462
I584	TTCGTACACAA ATCATGCAAA TTATCCTCAGT TACTTGCACA	e63	e1233	e250	e1272
A585	TGCGATATTT TACGTTGAGT TGTGTACGAA TTCGGTATAGA	e33	e740	e63	e243
I586	TACACGAACA TCCAAGACTAT AAAATATCGCA AAGCCATCTT	e669	e1237	e33	e1276
A587	ATTGCTGATT ATCATTGGGT TGTTCGTGA TTACTGGGTA	e573	e1239	e669	e406
A588	AGTTGTAGTCT AGCAGAAGAA TTGAGCTGAA AATGTCAGGA	e695	e568	e575	e320
A589	AGGGATGTAAT TTAGCTTGGT TTTTTTTTTT TTGTTGCATAA	e88	e359	nullT10N	e464
I590	TTACACCGAT TTTGCATCAAT ATTACATCCCT TTGCACCTTA	e31	e1257	e88	e404
A591	TTAGGTAGCAT AAAGGTGCGAT ATCGGTAA TGAGTAGAAC	e324	e466	e31	e578
I592	TTTGATTGCGAT TTGAGATCGT TCGAAGTTGT AGGTTCTAGTT	e332	e551	e326	e145
A593	TAGAGGAAAGT TGGTCTTGA TCGTTATGGT TACATGGAAGA	e153	e253	e334	e447
I594	TTGACACCAT TGTTCACTTT TTTTTTTTTT TCACCAAAC	e82	e370	nullT11N	e32
A595	TGTATGAAGGA TGTGCAGATA ATGGTGTCAA TAAGAAGGTCA	e287	e1272	e82	e231
A596	AGAGAAATGAGT AAGATGCCCT TCTTGTGTGT TATGACTGTGA	e450	e1276	e289	e378
A597	ATGGATTAGGA AGGCTGAAT TCTTGGGAA ATGATTGCAAT	e479	e576	e187	e198
A598	TTTTTTTTTT TTGGGTTCT AAGATGCTGA ACTAAAGGAGT	nullT11S	e697	e481	e1313
A599	AACTGATAGGA TTGGGTTAA TTATGCTGGT ACAATGGAAAA	e708	e327	e624	e239
I600	TGGATGATACT AGTTGATGCT TGTGTACGAA TATACGAAGGT	e41	e335	e63	e817
I601	TTTTTTTTTTT AGGGTGAAT ACGAGAAAAGT ATTTGCGAGAT	nullT11S	e509	e493	e841
I602	AGAGATGACTT AGTTGGTGA TTTTTTTTTT TATCGGAGAAA	e275	e32	nullT10N	e753
I603	TTGGTTTGAAT TGTGTGAAC AAATGGGTGA ACAATTGGAAA	e422	e290	e277	e480
A604	TTGACATTGTT ATGATGGGT TTATGGACGT ATGAGTTAGGT	e1341	e204	e244	e476
I605	TTTTTTTTTT ACTCCTTGTAA ACAATGTCAA TAACACCCAA	nullT10S	e1313	e1341	e150
A606	ATGTGAGTAGT TTGAGCAGAA TTAGTCGTGT TACATGATGGA	e283	e408	e436	e10
A607	AGGATCTGATT AAATGAGGCT TGTGCAGATAT ATGGTAGATCA	e1352	e347	e285	e742
I608	ACCCTTTGAT ACCTTCGTATA AATCAGATCCT ACTAGCACAA	e726	e817	e1352	e19
A609	ACTCTAGGGAA AAGACGGAAA TCTTGGGAA TAGAGGAACAT	e857	e265	e187	e789
A610	AGGAGTTACTT TTAGCGATGT TGACTGGAAA AAAACATTGAT	e381	e425	e859	e390
A611	AGGGTTAGAAT TAGTGGCAAT TAGAAAGCGA AGACTTAGGAA	e520	e87	e383	e126
A612	AAGGGATTAGT AGAGTGCCTT ATGACATGGT AAAGAGAGGAA	e37	e506	e782	e222
A613	TAGAAGGCATA TGTGAGATT TTGGAGTTT AAAATTAGCGT	e307	e286	e39	e81
A614	TGGTCATAGAT TTGTGCTAGT TGTACGGAAT AGGTTCTAGTT	e1384	e19	e771	e145

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
I615	ATGTCCACTT TCTTCCTTCT ATCTATGACCA ATGCCACTTA	e91	e729	e1384	e1387
A616	TTTTTTTTTT TACGGTGAAT AAGTGGACAT AGAATATGGGT	nullT11S	e90	e91	e713
A617	TTATCTGTGGA TAGTGAACGT TTTTTTTTT AAAAGTGAACA	e125	e843	nullT10N	e370
A618	TAGCATGAGTA AGTGTACGAA TATTGGTCGT AGCTAGTGTAA	e434	e763	e648	e34
A619	TGGGAATATCT TTTTTTTTT TATTGCGGT TTAGAGTGACA	e1	nullT10W	e203	e631
A620	AGTTTGTGTTGT TTTTTTTTT TCGAGAAAGTT ATGTAGTAGCA	e209	nullT10W	e3	e259
A621	AGATGTGATCT TTTTTTTTT TGGGAAGTTT TATATGACGGA	e197	nullT10W	e211	e210
I622	TGTAGATGAGT TGGGTCCTGA AATGTGAGGT AAAAATGGACA	e266	e253	e67	e394
A623	AGAATTAGGGA TGTGGTACAA TGGTTACGAT TCTAAGTGTGA	e108	e40	e268	e173
A624	TGGAAGAGTAT TAAGTGGCAT ATCTGGGTTT TTTAAATGCGA	e101	e1387	e174	e190
A625	TTGAGGGATAT AGTTGATGCT AAGGGAACAA TCAAGAAGGTA	e61	e335	e416	e42
A626	ACGAGATAGAA ATTGGGTCAT TGTGTACGAA ACAATTGGAAA	e140	e224	e63	e480
I627	AGGTATCGTAT TGTGCGTTAGT AGAACAGCCTT TTTCATTGGT	e315	e156	e142	e329
A628	TGTAGAGGATT TGTTGGTCTT AATGTCAGGT TGATTAGGAGT	e655	e380	e256	e414
A629	ATGTAGGGTAA TTTGGGTGAT AAATTGCGTT AGGATATAGCA	e303	e4	e534	e94
A630	TAGTGGTTGTA AATCGGAGTT TTGTGTAGCA ATAGTGAGGAA	e438	e200	e799	e325
A631	AAAAGGGTCTA ATGCTGAGAT AGGCAGTAAA AGAGTTAGGAA	e614	e269	e440	e194
A632	AATAGGGTACA TGTTACGAGT AGGTGCTTAA TCTGGGATTAT	e93	e111	e616	e235
A633	TGTAGATGAGT TTTGGGTGAT TCGAAGTTGT TAGGTTTCAGA	e266	e4	e326	e853
A634	TTTTTTTTTT TTTGGGTTCT AATGACGGTA AGAACAGGAT	nullT11S	e751	e460	e368
A635	ATGGGATTAGA TTCGGAAGAT TTTTTTTTT AGAGTGGATAA	e69	e417	nullT10N	e798
A636	TATGACAGAGT ATGATGAGCA ATTGAAGGGT ATATAGAGGCA	e524	e64	e71	e427
A637	ACAGTAGGATT TTGGATTGGA TGTTGATCGA TTTAGGACGTA	e157	e143	e526	e1492
A638	TAGCTAATGGA TCGGTTATGT ATGAGGTGTT ATATAGGACGT	e458	e457	e375	e496
A639	TTGGGATATCT ATGGAAGTGT AATGACGGTA AGGATATAGCA	e487	e658	e460	e94
A640	AAGTACTGAGA TAAGGCATGA TGTTAGCGAT ATTGTTTGACA	e121	e294	e867	e38
A641	TCGAGAGATTA TTTGGATGGT TTATGGCAGT AAGAGAGTGT	e405	e306	e123	e304
A642	AGAGTTTAGGA ACGTGTATGT AAAGATGGCT ATCAAGGTAGA	e168	e441	e371	e1516
A643	ACTTGTAGGAT AGCGTTGATA TTTGGGTACA AGGTTTAGGTA	e279	e36	e293	e639
A644	AGTCGATGATA TGTTGGAAC TGTGCGAGTTA TAAAAGGCAA	e475	e502	e281	e6
I645	TTTTTTTTTT ATCCTGATTCT AGTTTACTCCT AGAAACCCAA	nullT10S	e368	e364	e697
A646	AGGGTTCTAAAT ACTGGTGATT TCGCAGAAATA ATAGCAGGATA	e161	e72	e827	e423
A647	AGTGTGAGATA TTGAGCGATA AGTGAGTTGT TTAGAGTGTCT	e1536	e527	e473	e746
I648	ACGACACATA TACGTCCTAAA TATCTCACACT TGCCATATCA	e1538	e1492	e1536	e376
A649	TTATCTGTGGA TGATATGGCA TATGTGTCGT AGATCAGTGT	e125	e376	e1538	e169
A650	TAGATGTACGT TTGGATTGGA TCAAGTTGGT ATGGGATCATA	e630	e143	e542	e149
A651	TAAAATTGGCA TACGTTGTGA TCGTGGTATT AGAGTAGTGT	e319	e282	e632	e226
A652	TGGTAAGTACT TGGTATGCA TTTGTAGGCT TATACGAAGGT	e348	e124	e114	e817
A653	TAGGAACAGAA TTGAGCAGAA TTGGAACGTA ATGACAGTAGT	e344	e408	e127	e785
I654	AACATACCGA TCTACCTGT ATTAGAACCCCT ATTACATCCGT	e1562	e1516	e161	e1563
A655	TAGGAACAGAA TAAGGCATGA TCGGTATGTT AGCAAAAGAAA	e344	e294	e1562	e308
I656	TGCGATATTTT TACGTTGTGA TTATGCTGGT TACATGATGGA	e33	e282	e624	e10
A657	TGCGATATTTT ATTGCGTTTT TCAGGGAAAA TTAAAGAGGGA	e33	e815	e35	e566
I658	AACTCCCTTT TACTACCAACT AAAATATCGCA TACCCACAAA	e1574	e365	e33	e1575
A659	TTTTTTTTTT TTGGGTTCT AAAGGGAGTT AGTGAACAAAA	nullT11S	e697	e1574	e165
A660	TGAGTATCTGT AGTATGCTGT TTTTTTTTTT TTCATGTTGTT	e467	e828	nullT10N	e2
A661	AGAATTAGGGA TCGGTTATGT TGGTGCATAT ATTAGTGGTA	e108	e457	e223	e49
I662	TTCTATGTGGA TGATATGGCA TCGAGAAAGTT AGAACATATGGT	e213	e376	e3	e154
A663	TATCGGAAGTA TACGTTGGAT AATGGTGGAT TTTTAAGCGT	e176	e543	e215	e70
A664	AGGAAGGAATA TGGACAGTT TAATGGAGCA ATAGGTGACAT	e270	e515	e178	e1594
A665	TTTGACTGTTT TCTGTTGGT TCGTGAAGA ATTGTAGGACT	e844	e449	e272	e676
I666	AGTAGATACGT TGAAAGTCGA ATCGTAGAA TCAGTTAGGA	e503	e351	e58	e810
I667	ACAGTAGGATT AAATGAGGCT AATCAGTGGT ACTTTAGAGGT	e157	e347	e814	e180
A668	AGAAGTCAGAT ACGGATGAAT AGCTGTGATT TCAGTTAGGA	e352	e1563	e159	e810
A669	AGGTAAGAGAA AGTCAGAAT ACGTGTGTAT ACTAGATGGAA	e820	e686	e354	e647
A670	ATGAAAATGCT AGCGTTGATA TGATTGAGCT AAAATGCAAGA	e610	e36	e822	e316
A671	ACATATGGAGT TTTGTGGGTA TAGCAGGTTT TTTGGTTCAA	e581	e1575	e342	e113
I672	TTTTTTTTTT TTTTGTCACT ACTCCATATGT ATTACCGGTA	nullT10S	e165	e581	e90
A673	TGAGGACTATT TCGAGTGAAT ATCGGTGAA TGAGGTATGAT	e76	e396	e31	e45

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
A674	AGATGGAGATT TGTCTGTGTT TGGATATGCA TATAGACGTGT	e852	e776	e78	e255
A675	TGGTAGTAACT AGAGCAATGA AAGCGAGTTA AGGCTTGATA	e97	e216	e854	e821
I676	ATCGACATCA ATGTCACCTAT TCCTATCAGTT TTTTTTTTTT	e518	e1594	e708	nullT10E
A677	AGTTCTGAGAT AATGAGAGCT TGATGTCGAT TTTTTTTTTT	e717	e273	e518	nullT11E
A678	ATGGATCTGAT TTGGCTTGTA ATGGATGTGT TTTTTTTTTT	e890	e847	e718	nullT11E
A679	AGGTATGGAAT TTTGATCGGA TGCTTGGTA TTTTTTTTTT	e837	e160	e891	nullT11E
I680	TTGGGATATCT AGGTTAACGT AAAGATGGCT TGAAAGAGGTA	e487	e445	e371	e357
A681	TAGGAAAGTCAGGTTAAGGTTGAT TTGGCAGTAT TTATGTTGCA	e552	e613	e489	e410
I682	TTTTTTTTT TACGGTGAAT ACGGTTGTTA AGAACAGGAT	nullT11S	e90	e554	e368
I683	ACATATGGAGT AGTTTGGTA TTTTTTTTTT AACTATGAGGA	e581	e32	nullT10N	e137
A684	ATGGATGTCTA TGGGTGCTAA TCTTGGGTT TAAGTGACTGA	e299	e79	e747	e725
I685	TGAGTATCTGT TACGGTGGAT TCTGTTGGAA TGGAGTAAGAT	e467	e543	e301	e186
A686	AGTAGACTGAA TAAGCGTGTAGATGCGAA ACTGAGTAAGA	e73	e100	e163	e594
A687	TAACTGGAAGA AGCTGAGAAAT TTTTTTTTTT TCGAGAGATAA	e393	e840	nullT10N	e62
A688	AGTAATCTGGT ATGGACTTGT TCATGGATGT TAATAGTCGGA	e401	e694	e151	e251
A689	AGTGTGGAA TGTGACAAGA AAGCTGTGAT AGAGTTGAGTA	e454	e490	e403	e643
A690	TATATGGTCGT ATTGCGTTT TGTTCTGTGT ATTATCGGAGA	e193	e44	e166	e627
A691	AGACTAATGGA ATGGGTTGAA TCATGGGTTT ATAGAGTTGGA	e528	e302	e514	e98
A692	AGAAGGATGAT TTGTGCTAGT AAGTGGTGA TGGGTACTTTA	e638	e19	e510	e688
I693	TTTTTTTTT TGAGTGCATA TAGCAGGTT TGAGGTATGAT	nullT11S	e75	e342	e45
A694	ATGTAGGACTT TCGAGTGAAT ATGTCGGTAA TTTCATAGGGA	e295	e396	e106	e21
A695	ACAAGGATAGA TAAGGTGCAA TGTGACATGT TAAAGGGAGAA	e116	e404	e297	e549
I696	TGGCTTTTTTA TCGGTTATGT AAAGTGGTCT AAATTGGAACA	e512	e457	e620	e102
A697	TTTTTTTTT AGATGACGTT TCATGGGTT TAGGAAGACAA	nullT11S	e415	e514	e60
A698	TGGAGATACAT AACAGGTGAA ATCAAAGGGT AGCTAGTGTAA	e172	e641	e726	e34
I699	TTTTTTTTT ATCATAACCTCA ATGTATCTCCA TCCAATCACA	nullT10S	e45	e172	e1737
A700	TGCAGTTTTTA TTCGATGGAT TTTTTTTTT TCGAGAGATAA	e48	e107	nullT10N	e62
A701	TGGCTTTTTTA ATGGTGTGAA TAGCGATGAA TAAAATGTGCA	e512	e298	e50	e602
A702	ATACTGGATGAA TTGTAGCGAT TAATGGCTGT AGATTATCGGA	e129	e494	e46	e214
I703	AGGGATGTAAT TGGACAGTTT TTTCGAGGTT ATAGGACATGAA	e88	e515	e131	e525
I704	ACCCTAAAAA TAGCCTCAATA TCCTTAGACT TTACCAACAA	e1757	e541	e532	e327
A705	AGAACTAAGGA AAGCAGATGT TTTTGAGGTT TATTGAGGTA	e136	e128	e1757	e834
A706	TTTTTTTTT TGTGATTGGA TAGATGCGAA AGGTCTTGTAGT	nullT11S	e1737	e163	e736
A707	TTTAGTGGAGA AGGTTGACT AAGTGGACAT TAGGGTAACAT	e646	e51	e91	e117
A708	AACGTGAAATT AACAGGTGAA TATTGGTCGT TAGAGAATCGA	e258	e596	e648	e109
A709	AGGGATTCTATA AAGAAAGGCT AAGGCTGATT AAGAAAAGCAA	e792	e27	e260	e590
A710	TTTTTTTTT TACGGTGAAT AGCAAGAGAA ATCAAGGATGTA	nullT11S	e90	e734	e556
A711	ACTATGTGAGT TTGGCTATGA TTTTTTTTTT AGAGTAAGGTT	e430	e167	nullT10N	e858
I712	ACTTACTGCA TACCTCGAATA AATCCTACTGT TCTTCAGCAT	e1789	e834	e157	e1790
A713	AGTGATTAGGA ATTGGGTCAT TGCAGTAAGT TTTCATTGGT	e144	e224	e1789	e329
A714	TTTTTGTCGT TTTGATCGGA TGATGAAGCT AGATCTGAGTT	e764	e160	e146	e419
A715	ATTAATTGCGA ATTGGATCGA TTTTTTTTTT AGGCTAAGTAT	e516	e92	nullT10N	e873
A716	AGGTATCGTAT AAGGTACGTT TGATGTCGAT TTCTAGGAAGT	e315	e649	e518	e1803
A717	AGTAGACTGAA TGCTGTGTTA ATCGGATTGT TTTTTTTTTT	e73	e433	e887	nullT11E
A718	TTGCGATAAAA ATGCTGAAGA TTCGTGGTAA TTTTTTTTTT	e1816	e1790	e893	nullT11E
I719	TTACACGAA ACCAAATGAAA TTTTATCGCAA TTTTTTTTTT	e893	e329	e1816	nullT10E
A720	AAAGAGGAGAT AGTAGTGCAT TTGCTGGTAA TTTTTTTTTT	e719	e147	e893	nullT11E
A721	TTTTTTTTT TGCAGAAGAT TTTCGATGGA TTTTTTTTTT	nullT11S	e135	e720	nullT11E
A722	AGGAAAGTATT TGTGTCAGA AACTGTGTGA TTTTTTTTTT	e1821	e519	e716	nullT11E
A723	TGCCCTACAA ACTTCCTAGAA AATACTCCCT TTTTTTTTTT	e1822	e1803	e1821	nullT10E
A724	TGAGGATGTTA AGCAGGATT TTGTAAGGCA TTTTTTTTTT	e715	e339	e1822	nullT11E
A725	ATCACCGAAT TTCTCCCTAA TGTATCCTAA TTTTTTTTTT	e1826	e549	e185	nullT10E
A726	TTTTTTTTT AGTTGATCGT ATTCCGGTAT TTTTTTTTTT	nullT11S	e603	e1826	nullT11E
M727	AGATGTGATCT TTTTTTTTTT TTTTTTTTT TGGGTGTATAT	e197	nullT10W	nullT10N	e652
M728	AAGATACGAGA TTTTTTTTTT AATGAGGTT ATTGTTGACA	e20	nullT10W	e67	e38
M729	AGGTTAAGAGA TTTTTTTTTT AGGACTGTT ATAGAGTTGGA	e5	nullT10W	e939	e98
M730	TACGGTATTGA TTTTTTTTTT ACGTATGTCGAA TAAATGTCGAA	e16	nullT10W	e7	e602
M731	AGTTTGTGTTGTT TTTTTTTTTT TGTGATGCTA ATTACGAGAGA	e209	nullT10W	e18	e431
M732	TTAGGAAATCT TTTTTTTTTT TGGGAAGTTT ATAGCAGGATA	e917	nullT10W	e211	e423

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
M733	AGATGTCAGAT TTTTTTTTT TAGAGCGAAA AGAGTAGTGT	e913	nullT10W	e22	e226
M734	TGGGAATATCT TTTTTTTTT AGGTAAACGTT AGGTTAGACAT	e1	nullT10W	e26	e341
M735	TTTCGGATAGA TTTTTTTTT TCGAGAAAGTT TAATAGTCGGA	e28	nullT10W	e3	e251
M736	TGGCTATTAGA AATCGGAGTT ACGAGATTGA AGTGAACAAAA	e65	e200	e698	e165
M737	TTTAGTGGAGA TGGATCGATT TTGGAACGTA ATAGACAGGAA	e646	e8	e127	e57
M738	TGAGTATCTGT TTGTGGTGT TAATGGCTGT AGAACTATGGT	e467	e59	e46	e154
M739	TGGTAGTAACT TGCTGTGTTA TGGTGCTAT AATTTTCGGT	e97	e433	e223	e202
M740	AGGAAGGAATA TGGGTTCTT AGCTGAAGTT AAGGTAGTCAT	e270	e23	e99	e504
M741	AGAGTTTAGGA TCGAATGGA TCGTGAAGA ACAGGAAGATA	e168	e916	e272	e345
M742	TATGGAGTTCT AAGAAAGGCT TTTGGGTACA ATATAGGACGT	e385	e27	e293	e496
M743	TAGGAACAGAA TTTGGGTGAT ATAACGTGGA TGTGACAGATA	e344	e4	e387	e778
M744	AGGACTAATGT TAGGCAATGA AGCTGTGATT AGAGAAGATGT	e659	e196	e159	e619
M745	TTTTTTTTTT TGTGTTGAGT TCTTGTGTT ATGCCAAGATT	nullT11S	e30	e661	e120
M746	AATAGGGTACA TGAAATGGGT ATGGTTGTCA TGGTATTTCAGA	e93	e68	e195	e85
M747	AAGTTAGGGTA AAGCAGATGT AAGGGTCAAT TAATACGAGGA	e84	e128	e95	e74
M748	ACTTGTAGGAT AAGGTACGTT AAAGAAGGGT ATATAGAGGCA	e279	e649	e86	e427
M749	AGATGGAGATT TTCGGAGATT TGTGCAGTTA TATACGAAGGT	e852	e478	e281	e817
M750	TGTTGTTGATT TCGAGTGAAT AAGCGAGTTA AGAAAAGGAAGA	e246	e396	e854	e729
M751	TAAGAGTGTGA TAAGCGTGT AGTTCGGATT TAAGAAGGTCA	e548	e100	e248	e231
M752	ACGTTGATTTT AATGAGAGCT TGAGACGAAA TATTGAGGCTA	e565	e273	e428	e541
M753	TTATCTGTGGA TAAGGCATGA TTAGCGAGTT TTCTGTAGTGA	e125	e294	e567	e398
M754	ATTGCTGATT AGTTCTGGTT TCAAGTTGGT AAGAAAAGCAA	e573	e388	e542	e590
M755	TGGACAAATTT AAGGAGGTTT TAAAGGTGCT AGATCTGAGTT	e148	e179	e665	e419
M756	TGGAAGAGTAT TGGCTGTAAT TTTGTTAGGCT TCAAGAAGGTA	e101	e96	e114	e42
M757	AGGGTTCTAAT TAGTGGCAAT AGACTGGTTT TTTAAATGCGA	e161	e87	e103	e190
M758	AGGATCTGATT TACGTTGTGA AGTGAGTTGT TTAAGGGAGAA	e1352	e282	e473	e549
M759	TGGTCATAGAT TACGTTGGAT ATCAAAGGGT AGTATTACGGA	e1384	e543	e726	e53
M760	AAGTTAGGGTA AGCAGAAAGAA AAGACCAAGTT AGAAAGTAGACA	e84	e568	e587	e386
M761	AGAGTTTAGGA TGGACTTTGT TGACGTTGTA AAAGGCTAGTA	e168	e808	e546	e288
M762	AGAGTGTATGAT TGAACGTGAA TGATGAAGCT ACAATGGAAAA	e463	e666	e146	e239
M763	TTTTTTTTTT TTGGGTGTTA AGGAACAAAGT TGAGGATTACA	nullT11S	e150	e54	e2014
M764	TGTTTGAACCTT TAGTGAACGT TTTTTTTTT TAGTTAGTCGT	e133	e843	nullT10N	e276
M765	ATGGGATTAGA AGCAAGGATT AGGTGAATGT TAGAGAATCGA	e69	e115	e538	e109
M766	TCGTAGTATGT TGGCAGTAAT ATTGAAGGGT AAATTGGAACA	e618	e104	e71	e102
M767	TAGATGTACGT TAAGTGGCAT ATGGTGTCAA AATGTCTAGGA	e630	e1387	e82	e320
M768	TGGAGATACAT TGTGGAACAT TCGTGGTATT ATTGATGCAAA	e172	e502	e632	e1257
M769	ACTATGTGAGT TTGAGATCGT ATCTGGTTT ATAGTCTGG	e430	e551	e174	e1237
M770	TTTGATTTCGT TTAGCTTGGT TGTAGCGAT ATGGTAGATCA	e332	e359	e867	e742
M771	ATGTGAGTAGT TGGTTTGCTA TCGTTATGGT AGGAAATAGGA	e283	e171	e334	e757
M772	TGAGATTGGTA AAAGGTCGAT TGTGGATAT AAGAGAGTGT	e2059	e466	e285	e304
M773	TTTTTTTTTT TGTAATCCTCA TACCAATCTCA ATCACATCGT	nullT10S	e2014	e2059	e2061
M774	ATACTGGATGA ACTGGTGATT AGGAACAAAGT AAAAATGGACA	e129	e72	e54	e394
M775	ATTGCAATGAT AGCTTGTGTA TTTCGAGGTT TGAGGTATGAT	e700	e621	e131	e45
M776	ATGGTTTTGTT AACAGGTTGA AGGGATAGGA AGAAAAGGAAGA	e356	e596	e702	e729
M777	AGGAGTAAACT TGGTGGTAA TGTGATACGT AAAGAGAGGAA	e364	e83	e358	e222
M778	AGGTGAAGTAT AGTTGTGCT TGTACGGAAT TGAGTAGAAC	e784	e335	e771	e578
M779	TGGAAGTACT TGTGAGGATT TTAGAGCGAA AGTTGGTAGTA	e348	e286	e309	e365
M780	TTTTTTTTTT ACGATGTGAT TTGGAACGTA TAGGAAGACAA	nullT11S	e2061	e127	e60
M781	TCGGAATTAGA ATAACGGTGA TTTTTTTTT TTGTTGCATAA	e597	e323	nullT10N	e464
M782	TAGTGGTTGTA AGCAGGATT TAGAATGCGA TTAGAGTGTCT	e438	e339	e628	e746
M783	TATGGAGCTAT TTGAGTGTGT TCTTGGTTT TGTAGGAATCT	e756	e703	e747	e374
M784	AGGGTTCTAAT TAAGGTGCAA ACGTGAAGA AAAAGTTGTGT	e161	e404	e43	e206
M785	TTCGAAATGTT AAGATGGCTT TAGATGCGAA AGAAGTAGACA	e557	e1276	e163	e386
M786	AGGGATTAGT TTGTGGCTTA TGTTCAAGGTT ATAGGTGACAT	e389	e429	e559	e1594
M787	AGAGAATGACT TTGGTCAGAA TATGTTTCGT ATTATCAGGGT	e450	e184	e391	e158
M788	AGGGATGTAAT TGAAAGTCGA AAAGGGAGTT AATGTGGAAAA	e88	e351	e1574	e141
M789	AGAATTAGGAA ATGGGAACCTT AAGTGGTGAA TTAAGGAAGGT	e108	e629	e510	e806
M790	ACATATGGAGT ACGTGTATGT TCGAGAAAGTT TAAGTGAECTGA	e581	e441	e3	e725
M791	TTAAGGACTGA TGGTATGCAT TTCTGGTAA TTTTTTTTTT	e712	e124	e893	nullT11E

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
M792	ACTGAGGATAA TGTGTCAAGT ATGTGGATCA TTTTCGATTGT	e250	e572	e758	e845
M793	AACTGATAGGA AGTCGGATT AATGGTGGAT TTTTTTTTTT	e708	e560	e215	nullT11E
M794	AGTGGATCTTA TTTGTGGGT AAGCAAGAGAA AATAGTCTGGT	e777	e1575	e734	e130
M795	AACAGAGAGAT TTGGCTATGA TTTTTTTTT AGAGTGGATAA	e833	e167	nullT10N	e798
M796	TATATGGTCGT TGTCTGTGTT ATGGTCAAAT TTTCATAGGA	e193	e776	e835	e21
M797	TCACCCATTT TATCTTCCTGT TTTTTTTTTT TACCACAACA	e277	e345	nullT11N	e2209
M798	TGGCTATTAGA AAGGTTCGAT TTTTTTTTTT TTAGTCTTGGA	e65	e188	nullT10N	e2220
M799	AGTGATAGGT ATCGAGTGT TAGTTGCGAT TCTTAGAAGGT	e189	e735	e730	e218
M800	AGGACTAACT AGTGGAAATGT TCGGTGAAA ATAGGACATGA	e364	e780	e191	e525
M801	AGGGATTCTA ATGTATGGCT TCGGAGAATA TTAAAGAGGGA	e792	e836	e827	e566
M802	AGGAAGATTCT TTGTAGCGAT AGGCAGTAAA AGGTTAGGTA	e1076	e494	e440	e639
M803	TATGACAGAGT TGTTGTGGTA TTTTTTTTT ACTTGGTAGAT	e524	e2209	nullT10N	e1184
M804	AGGATGTACTT TTAGCTTGTG TGTTGATCGA TTAGTAGGTGT	e829	e359	e526	e29
M805	TGTGGAGATAT TGTGTCTTGA TAAAGCTGGT TAGTTGTGAGA	e2249	e318	e831	e451
M806	TTTTTTTTT TCCATAACTCT ATATCTCCACA TCACTTGTCA	nullT10S	e462	e2249	e2251
M807	AACAACCTGA TCCAAGACTAA TTTTTTTTTT TTACAGCCTT	e2252	e2220	nullT11N	e2253
M808	TAGTCAGTTGA ATTGGACGAT TCGAGTTGTT TAGGTTTCAGA	e262	e731	e2252	e853
M809	ATGAAAATGCT AGTATGCTGT TTTTTTTTT TCGAGAGATAA	e610	e828	nullT10N	e62
M810	TTTTTTTTT ATCTGCAAAAT TATCATCGACT ATCACACGAT	nullT10S	e841	e475	e2273
M811	AGTAGTGAAGT TTGAGCGATA TGAAGAGCTT AATCGGAAAAAA	e56	e527	e1230	e1036
M812	TTTTTTTTTT TGACAAGTGA TTAGTCGTGT ACTAAAGGAGT	nullT11S	e2251	e436	e1313
M813	AGTCGATGATA AAGGCTGTAA TTTTTTTTTT TTCATGTTGTT	e475	e2253	nullT10N	e2
M814	AAGAGGTAGAA AAGACGAAA TCTTGTGTGT ATGTAGTAGCA	e1236	e265	e289	e259
M815	TGATAGGACTT TGTTGAACT AGACATGGTT TGGAGACTAAT	e884	e502	e1238	e296
M816	AGTAGTGCACAT AGGTTGTGA AAGTGGACAT AGTAATCAGGA	e589	e613	e91	e983
M817	AGTCTAAAGGA AGTGCAGAAT TTTGAGTGGTA TTTCATTTGGT	e532	e686	e591	e329
M818	TTTTTTTTT ATCGTGTGAT AAATTGCGTT AGAGTTATGGA	nullT11S	e2273	e534	e462
M819	AGAAGTACAGA ATAACCGTGA TTTTTTTTTT ATTCAAGTAGT	e1260	e323	nullT10N	e2308
M820	AGGTATGGAAT TTGTGGTAT TTCAAAAGGT TGAGAGAAAGA	e837	e59	e1262	e733
M821	TTGACATTGTT AGAGTGTCTT TGATGATCGT TTCTGTAGTGA	e1341	e506	e313	e398
M822	ATGTAGGGTAA ATTTGCGTTT ATCGGTGAA TTTAAATGCGA	e303	e815	e31	e190
M823	TTAGGTAGCAT ATCATTGGGT AAAGAAGGGT TAAGAAGGTCA	e324	e1239	e86	e231
M824	TTTTTTGCGT ATTGGATCGA TTTTTTTTTT AGTGTAAAGAGA	e1027	e92	nullT10N	e738
M825	TAGAGGAAAGT TTGTGCTAGT ATGAGGTGTT AGTGGTAGTA	e153	e19	e375	e365
M826	ACCTTCAGTT ACTACTCGAA TTTTTTTTTT AACAAATCCCT	e653	e2308	nullT11N	e2341
M827	TTCGGATAGAA AGCTGAGAAT AGAACAGCGT AGAGTTAGGAA	e651	e840	e142	e194
M828	TTTTTTTTTT TTGGGTGTT AACTGAAGGT ATTTTGACGT	nullT11S	e150	e653	e105
M829	AGAACAGAGTA AGTTTGTGA TTTTTTTTTT AGTAGGAAGAA	e234	e32	nullT10N	e122
M830	AGAGATGACTT TTGGATGTT TGGATGTCAA ATGAGTTAGGT	e275	e306	e236	e476
M831	AGTAGACTGAA AGAGCGTAAT AAATGGTGA TAGAGAATCGA	e73	e343	e277	e109
M832	TTTTTTTTTT AGGTTGGAAT TGAGGCAAT AGAACATCAGGAT	nullT11S	e509	e411	e368
M833	AATGCAGATT AGGGATTGTT TTTTTTTTTT ATCGGTAGTAA	e328	e2341	nullT10N	e2374
M834	TTGAGAGTAGT AGTGTGCAAA TTTGGAGTGT ACAATTGGAAA	e238	e763	e330	e480
M835	AGGGTTAGAAT TTAGGCAAGT TCATGGATGT AGATTATCGGA	e520	e237	e151	e214
M836	AGAACCTAAGGA AAGACGAAA TAATGGAGCA AGTACGAGTAT	e136	e265	e178	e1049
M837	TTTTTTTTTT TGAGTGCATA TTGTTGGCTA AGAACATGGGT	nullT11S	e75	e138	e713
M838	AGAACCTAAGGA TGGAAAGAAA TTTTTTTTTT TGTTGATGTT	e136	e768	nullT10N	e2396
M839	AGCTTAGATGA TGTGGTCTT TGGTGCATAT ATTAGTCTGGA	e181	e380	e223	e623
M840	AGGAAGGAATA ATGAAGACGA AGAACAGAAGA AGATAGCGTAT	e270	e412	e183	e598
M841	AACAGAACCT TTACTACCGAT TTTTTTTTTT TCACCAAACCT	e2406	e2374	nullT11N	e32
M842	ATCTTAGGGAA AAGGGACAAT AGGTTCTGTT AAAATGCAAGA	e857	e331	e2406	e316
M843	AGGGATGTAAT ATGTAGCAGT TGACTGGAAA AGGATATAGCA	e88	e241	e859	e94
M844	TTTTTTTTTT TGCGAGAAGAT TTGTGTAGCA ACTTTAGAGGT	nullT11S	e135	e799	e180
M845	AACGTGAAATT AAGGGTGAAT TTTTTTTTTT AAAAGTGAACA	e258	e152	nullT10N	e370
M846	AGGTACGATTA TTTGGGTGAT AAGGCTGATT TAGGGATGAAA	e1093	e4	e260	e267
M847	TCAGCATCTT AACATCAAACA TTTTTTTTTT ACCCATGAAA	e481	e2396	nullT11N	e2429
M848	TAGAGATGCTT TTGGTCAGAA TGTCGTGTA TGCGATATAGGT	e1060	e184	e669	e705
M849	TTTTTTTTTT AGATGACGTT AAGGAATCGA ATTACAGAGGT	nullT11S	e415	e240	e796
M850	AAGTACTGAGA AGTGATCGAA TCTTGTGTGT TTAGTTCAGGA	e121	e763	e289	e263

Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
M851	AAGTCGAGTAT ATGATGGTGT TTATGGCAGT ATTAGTGGGTA	e225	e204	e123	e49
M852	TTTTTTTTTT TTTGGGTCT TTTGAGGCT TGATAGTGCTA	nullT11S	e751	e114	e2460
M853	TGGAGATACAT TTTCATGGGT TTTTTTTTT TTGTTGCATAA	e172	e2429	nullT10N	e464
M854	TCGGTAGAGTA TTTCTGGGA TATTGCGAGT AGAGTTGAGTA	e741	e670	e232	e643
M855	TAGAAGGCATA TGGCTGTTAA ATGTCGGTAA ATCAAGGTAGA	e307	e363	e106	e1516
M856	TTTTTTTTTT AGTTGGTCA TGGATATGCA ATAGGACAAGT	nullT11S	e227	e78	e650
M857	ATGGATTAGGA TTGTCAGGAA ACGATGGAAT TAGGGTAACAT	e479	e314	e1082	e117
M858	AGATAGTGGT TCTGTTGGT AAGATGCTGA AAAATTAGCGT	e2491	e449	e481	e81
M859	TTTTTTTTTT TAGCACTATCA AACCACTATCT TGCTAACCAT	nullT10S	e2460	e2491	e2493
M860	ACAGTAGGATT ATGATGGTGT AAAGTGGTCT TTTGTCATGT	e157	e204	e620	e963
M861	TTTTTTTTTT AGTCGTTGA TGCGATTAAGT ATTGAAGCAA	nullT11S	e826	e1789	e274
M862	AGGGTTCTAA TTCGATGGAT TTTTTTTTTT AGGAGTTACT	e161	e107	nullT10N	e17
M863	TTATCTGTGA TGTAAGTGTCT CGGTATGTT AGCTAGTGT	e125	e310	e1562	e34
M864	AGTTGAGTCT TGTTGCTAA TATTGGTCGT ATGATTGCAAT	e695	e79	e648	e198
M865	AACTCCAAC TCCCTCATAGTT TTTTTTTTTT TACCAATGCA	e2516	e137	nullT11N	e2517
M866	AAAGTGAGGATA TTGAGCGATA AGTTGGAGTT AGACTTAGGAA	e801	e527	e2516	e126
M867	TTTTTTTTTT ATGGTTAGCA AAGGTCAGT ATCAGGATGTA	nullT11S	e2493	e110	e556
M868	AGTCTAAAGGT TGGGTTCTGA TTTTTTTTTT AACTATGAGGA	e737	e253	nullT10N	e137
M869	ACAGGTTATTA TTGTAGCGAT AGTCGTCGAA TGGAGTAAGAT	e1007	e494	e739	e186
M870	TAAAATTGGCA ATGCTGAAGA TTATGCTGGT TATTGAGGTA	e319	e1790	e624	e834
M871	TTGAGGGATAT ACGGATGAAT TAATGGCTGT TAAAACGTGTT	e61	e1563	e46	e660
M872	TTTTTTTTTT TTGGGTTCT TTATGGACGT TTGACGAAAAT	nullT11S	e697	e244	e781
M873	TAGAGGAAAGT TGCGATGGTA TTTTTTTTTT AGGAGAATTCT	e153	e2517	nullT10N	e2550
M874	TAAGAGTGTGA TGTTACGAGT TGATGTCGAT TCAAGAAGGTA	e548	e111	e518	e42
M875	ATTGCTGATT TACGTTGAGT TGTTCTGTGT TATTGAGGCTA	e573	e740	e166	e541
M876	AGGAGTGTATT AAGAAAGTCGA TTGAGCTGAA TATAGACGTGT	e861	e139	e575	e255
M877	TTTTTTTTTT AGCTGATTGA ACGGAGAAAT AGGTCAATTAGT	nullT11S	e321	e863	e736
M878	AGGTAATGGTA ACGAGTTGAA TTTTTTTTT TAGCTGAGTAA	e704	e47	nullT10N	e2572
M879	TTTGACTGTTT ATGATGAGCA AGCTGAAGTT TTTACTGGGTA	e844	e64	e99	e406
M880	AGTATGTCGA TAGTGGCAAT ATGCGTAGAA TTAGAGTGACA	e809	e87	e58	e631
M881	TGCCAACTTA AGAATTCTCCT TTTTTTTTTT TAGCATCCAA	e2582	e2550	nullT11N	e2583
M882	AAGGGATTAGT TGTCGTTAGT TAAGTTGGCA AGGAGCATATA	e37	e156	e2582	e537
M883	AGGAGTTACTT TGTTGTCAGA TTCGGAGTT TAGGGTAACAT	e381	e519	e39	e117
M884	TTTTTTTTTT AGTTGATCGT TAGAAAGCGA TGAGGTATGAT	nullT11S	e603	e383	e45
M885	AGTCTAAAGGA TTGGCTATGA TTTTTTTTTT AGAGCTATGAT	e532	e167	nullT10N	e182
M886	ATGGATGTCATA AGGGCTAAAT TTTGAGGGT AAAGCATTGAT	e299	e576	e1757	e390
M887	TTTTGTCTGT AACTGTGGTA TCTGTTGGAA AGGTCTTGATA	e764	e864	e301	e821
M888	TGCTACAACT TTACTCAGCTA TTTTTTTTTT TCACCAAAC	e2604	e2572	nullT11N	e32
M889	AGAGAAATGAGT TTGGCAGTTA AGTTGTAGCA AGATAGCGTAT	e450	e707	e2604	e598
M890	AGACTAATCGA TTGGCTGTA CTTTGGAA TGTCATAGACA	e528	e847	e187	e25
M891	TTTTTTTTTT AGGTGCTTT AAATTGCGTT TGATTGTTGT	nullT11S	e811	e534	e2614
M892	TGAGGACTATT TTGGATGCTA TTTTTTTTTT AAAAGTGAACA	e76	e2583	nullT10N	e370
M893	AGTGATTAGGA TGTGGTACAA AGGCAGTAAA TAGAGGAACAT	e144	e40	e440	e789
M894	TAGCATGAGTA TTGGTGGTAA AAGGTCAGT ATGACAGTAGT	e434	e327	e110	e785
M895	TGAGTATCTGT ATGGGTGAA TTAGTCGTGT AGATCAGTGT	e467	e302	e436	e169
M896	TGGCAAAATAA AGGAACGATT AAAGAGGGTT AGGTTCTAGTT	e52	e12	e155	e145
M897	TTTTTTTTTT ACAACAAATCA ATGCTACCTAA ACACCTCAA	nullT10S	e2614	e324	e2647
M898	TGGGATCTATT TGTGTCATGA AAGGGAACAA TTTAGGACGTA	e816	e791	e416	e1492
M899	AAAAGGGTCTA AGTAGTGCAT ATGACGTGTA TTTTGCATGAT	e614	e147	e818	e1233
M900	TTTTTTTTTT TGTGATTGGA AGGTGCTAA ATTTGCAGAT	nullT11S	e1737	e616	e841
M901	AGTAGATACGT AGGAGGTAGA TTTTTTTTTT AGGCTAAGTAT	e503	e229	nullT10N	e873
M902	TTCTATGTGGA ATTGGGTCTA AATCAGTGGT ATAGTGAGGAA	e213	e224	e814	e325
M903	AACTGATAGGA ATGTGGTCTT ATGACATGGT TCTGGGATTAT	e708	e645	e782	e235
M904	AGACGGTAATA AAGAACTGGT TGTGTCGAA TGAAGTAAGGA	e606	e55	e63	e271
M905	TTTTTTTTTT TTGGAAGTGT TTGGACGTTAA TCAAAATGTGT	nullT11S	e2647	e608	e2680
M906	AGTGTGAGATA TTGGAAAGAT TTTTTTTTTT AGAGTAAGGTT	e1536	e417	nullT10N	e858
M907	ACTGAGGATAAA TTGAGTCGAT TATGTCGTGT AGATCTGAGTT	e250	e819	e1538	e419
M908	TTAAGGACTGA ATTTGCGTTT AACTGTGTGA TTTTTTTTTT	e712	e815	e716	nullT11E
M909	TTGCGATAAAA AGAGCAATGA AGTAGCGAAT TTTTTTTTTT	e1816	e216	e714	nullT11E


Name	Sequence (5' → 3')	Gl. 1/E	Gl. 2/N	Gl. 3/W	Gl. 4/S
M910	ATTCGTAGAGA TGGTTCTTGA TTTTTTTTTT TTTTTTTTTT	e897	e253	nullT10N	nullT11E
M911	TGAGGATGTTA AAATGAGGCT TGCAATGAGAT TTTTTTTTTT	e715	e347	e898	nullT11E
M912	AGGTTAACACT TTGAGATCGT AACTGTGTGA TTTTTTTTTT	e2701	e551	e716	nullT11E
M913	TTTTTTTTT ACACATTGTA AGCTTAACCT TTTTTTTTTT	nullT10S	e2680	e2701	nullT10E
M914	AAAGAGGAGAT TGATATGGCA ATCGGATTGT TTTTTTTTTT	e719	e376	e887	nullT11E
M915	TTAGGATGACA TGTGCAGATA TTTCGATGGA TTTTTTTTTT	e185	e1272	e720	nullT11E
M916	TTTTTTTTT TACGGTGAAT ATCGGATTGT TTTTTTTTTT	nullT11S	e90	e887	nullT11E

3.2 Fluorophore- and quencher-modified strands

The sequences below use Integrated DNA Technologies' (IDT) code format for DNA modifications.

Name	Tile	Sequence
H-ROX	H173	/56-ROXN/TTTGGAGTGT ACTTTAGAGGT TTTTGATTGCT TGTCGAGATT
	H148	/5IAbRQ/ATGTGAGTAGT TTAGGCAAGT TGTGCAGTTA AAATTGGAACA
H-FAM	H306	/56-FAM/TTTGAGTGGAGT GAGTGAAGA TGGCTTTTTA AGAACCGGATT
	H281	/5IABkFQ/ACAGGTTAGAT ATGGTTGGAT TCAAGTTGGT ACAGGAAGATA
H-five	H227	/5ATT0550N/TAGTCGTTGTA TGTAGTGCT TTAGTCGTTGTTTTAAAGCGT
	H252	/5IAbRQ/ACTGGGAATT TAATACGAGGA TTGGGATATCT ACGTGTATGT
H-six	H428	/5ATT0647NN/AGGTAAGAGAA TGTGTATGTA ATGACGTGTA AGGTCTTGATA
	H445	/5IAbRQ/TAGATGCGAA TAGGTTTCAGA AGATGGAGATT AGGTTAACGT
A-ROX	A636	/56-ROXN/TATGACAGAGT ATGATGAGCA ATTGAAGGGT ATATAGAGGCA
	A647	/5IAbRQ/AGTGAGTTGT TTAGAGTGTCT AGTGTGAGATA TTGAGCGATA
A-FAM	A544	/56-FAM/TCTTGGTTGGT ATATGGTAGCT AATGCAAGATTT TTGTGGCTTA
	A528	/5IABkFQ/TTTGATTGCT TGGTATGCAT ACGGAGAAAT AGTTGGTAGTA
A-five	A539	/5ATT0550N/TGGCTATTAGA TGGTTTGCTA TTGTTGGCTA TTTTCGATTGT
	A550	/5IAbRQ/AATGTGAGGT TGAAGTAAGGA AGGGATTTAGT TGAAATGGGT
A-six	A643	/5ATT0647NN/TTTGGGTACA AGGTTAGGTA ACTTGTAGGAT AGCCTTGATA
	A631	/5IAbRQ/AAAAGGGTCTA ATGCTGAGAT AGGCAGTAAA AGAGTTAGGAA
M-ROX	M863	/56-ROXN/TCGGTATGTT AGCTAGTGTAA TTATCTGTGGA TGTAAGTGCT
	M855	/5IAbRQ/TAGAAGGCATA TGGCTGTTAA ATGTCGGTAA ATCAAGGTAGA
M-FAM	M830	/56-FAM/AGAGATGACTT TTTGGATGGT TGGATGTCAA ATGAGTTAGGT
	M836	/5IABkFQ/TAATGGGCA AGTACGAGTAT AGAACTAAGGA AAGACGGAAA
M-five	M761	/5ATT0550N/TGACGTTGTA AAAGGCTAGTA AGAGTTTAGGA TGGACTTTGT
	M754	/5IAbRQ/ATTTGCTGATT AGTTCTGGTT TCAAGTTGGT AAGAAAAGCAA
M-six	M759	/5ATT0647NN/TGGTCATAGAT TACGTTGGAT ATCAAAGGGT AGTATTACGGA
	M767	/5IAbRQ/ATGGTGTCAA AATGCTAGGA TAGATGTACGT TAAGTGGCAT

3.3 Shape layouts



H: 480 tiles, 900 bonds, 175 unique tiles, 395 shared (168 all, 81 with A, 56 with M)

H0	S1	N2	S3	H4	N5	H6	H7			H8	S9	N10	S11	H12	S13	H14	S15
S16	B17	N18	H19	S20	N21	N22	H23			S24	B25	N26	H27	B28	H29	S30	H31
B32	S33	B34	S35	H36	S37	B38	S39			H40	S41	H42	S43	H44	S45	N46	S47
S48	B49	S50	H51	S52	H53	S54	H55			S56	H57	S58	H59	B60	B61	N62	H63
H64	S65	H66	S67	H68	S69	H70	B71			H72	S73	H74	N75	H76	S77	H78	B79
S80	H81	S82	H83	N84	N85	S86	B87			S88	H89	S90	B91	S92	H93	N94	N95
H96	B97	H98	B99	B100	B101	H102	B103	H104	S105	H106	N107	H108	S109	H110	S111	H112	S113
S120	N121	B122	H123	S124	H125	S126	H127	S128	H129	S130	N131	S132	B133	N134	B135	S136	H137
H144	S145	H146	S147	H148	S149	H150	N151	N152	S153	H154	N155	H156	S157	H158	S159	H160	S161
B168	N169	N170	H171	S172	H173	S174	H175	S176	B177	S178	B179	S180	H181	S182	N183	N184	H185
N192	N193	N194	S195	H196	S197	H198	S199	H200	B201	H202	S203	H204	N205	N206	N207	H208	N209
S216	H217	N218	H219	N220	H221	S222	H223	S224	H225	S226	H227	S228	N229	B230	H231	S232	N233
H240	S241	N242	S243	N244	S245	H246	S247	H248	S249	B250	N251	H252	N253	H254	N255	H256	N257
S264	H265	N266	B267	S268	B269	S270	H271	S272	N273	N274	N275	S276	H277	S278	N279	S280	H281
N288	S289	H290	S291	H292	S293	H294	B295	N296	S297	H298	B299	B300	S301	H302	S303	H304	S305
S312	H313	S314	H315	S316	H317	S318	H319	B320	H321	S322	H323	S324	H325	S326	H327	S328	H329
H336	S337	B338	N339	H340	S341	H342	S343	H344	S345	H346	N347	H348	S349	N350	N351	H352	S353
B360	H361	B362	H363	S364	N365	S366	H367	S368	H369	S370	H371	N372	H373	S374	H375	S376	H377
H384	S385	H386	B387	H388	S389	B390	S391									B392	S393
B400	H401	S402	B403	S404	H405	S406	H407									S394	S395
H416	S417	H418	B419	H420	S421	H422	B423									N424	S425
S432	H433	S434	H435	S436	H437	S438	H439									S440	H441
H448	S449	H450	S451	H452	S453	H454	S455									H456	S457
S464	N465	S466	H467	N468	H469	S470	B471									S472	N473

A: 496 tiles, 926 bonds, 181 unique tiles, 315 shared (168 all, 81 with H, 66 with M)

A480	A481	N10	S13	A482	N5	A483	I484	A485	I486	A487	S3	I488	S11	A489	A490	I491	S109	I492	I493	A494	S9	A495	S15
S120	I496	I497	I498	S301	A499	N351	A500	S345	I501	S303	A502	S126	A503	N115	A504	S213	A505	N251	A506	S305	A507	N282	A508
A509	S434	A510	S412	N229	S45	A511	I512	I513	S247	N308	S324	A514	N218	I515	S395	A516	I517	A518	S436	A519	S332	I520	I521
S80	N206	S50	A522	S188	N152	S90	I523	S54	A524	I525	I526	S278	A527	S451	A528	S222	A529	I530	A531	I532	A533	S382	A534
A535	S337	A536	I537	I538	S73	A539	S35	A540	S364	A541	N134	A542	S289	A543	S341	A544	S172	A545	N22	N296	S421	A546	S167
S24	A547	S343	A548	N207	A549	S457	A550	S203	A551	S410	A552	S406	A553	S138	A554	N62	A555	N309	A556	S124	A557	S314	A558
A559	S30	A560	S130	A561	I562	I563	N165	A564	S284	I565	S161	A566	S397	N169	S178	A567	S276	N350	S43	I568	N257	A569	S431
I570	A571	S385	A572	I573	A574	S293	A575	S476	A576	S466	I577	S474	N465	S368	A578	S186	A579	S241	A580	I581	A582	S132	A583
N192	I584	A585	I586	A587	S297	A588	S359									A589	I590	A591	N170	I592	S174	A593	S263
I594	A595	N151	A596	S234	A597	S249	A598									S312	N233	S322	A599	S366	I600	N255	I601
I602	S145	I603	N220	N21	S128	A604	I605									N424	S261	A606	S149	A607	I608	N379	S455
S440	A609	S449	A610	S199	A611	S270	N95									S408	A612	S20	A613	S402	A614	I615	A616
A617	S459	A618	S226	N262	S82	N116	S197	N2	N107	A619	S1	A620	S111	A621	S105	I622	N140	A623	S58	N121	S92	A624	S391
S216	A625	S33	A626	N75	I627	S291	A628	N339	N273	S349	A629	S159	N46	S417	A630	S228	A631	S318	A632	S259	A633	S376	A634
A635	S37	A636	S272	A637	S195	A638	N238	A639	S453	A640	S65	A641	N211	N131	N193	A642	S153	A643	S147	A644	S461	N189	I645
S432	A646	S245	A647	I648	A649	S280	A650	S326	A651	S232	A652	S182	A653	S180	N85	I654	A655	S353	I656	N18	A657	I658	A659
A660	S243	A661	S404	I662	S113	A663	N94	A664	S142	A665	S442	I666	S425	I667	N84	A668	N184	A669	N429	A670	S316	A671	I672
S16	A673	S41	A674	S446	A675	S52	N365	I676	A677	N372	A678	N478	N477	N468	A679	S438	S166	N357	I680	N253	A681	N286	I682
I683	S389	A684	S157	I685	S444	A686	S39									A687	N205	A688	N209	A689	N236	N141	S215
S88	A690	S393	A691	N274	A692	N330	I693									S56	A694	N155	A695	S355	I696	N266	A697
N288	N347	N279	S67	N183	N378	A698	I699									A700	N26	A701	S307	A702	S69	I703	S47
S264	N275	I704	A705	S117	N244	S86	A706									S48	A707	S334	A708	S136	A709	S414	A710
A711	S224	N194	I712	A713	S77	A714	S399									A715	S268	A716	S176	N242	S328	N283	S311
S464	A717	S470	A718	I719	A720	S374	A721									S472	A722	A723	A724	S370	N473	A725	A726

M: 480 tiles, 880 bonds, 190 unique tiles, 290 shared (168 all, 56 with H, 66 with A)

M727	S105	M728	I493	M729	S3	M730	S9	M731	S111	M732	I486	I492	S11	M733	I484	I488	S13	M734	S1	M735	S109	I491	S15
B360	M736	S35	B458	S67	M737	S334	B460	S247	M738	S243	M739	S52	M740	S142	M741	S153	M742	B201	M743	S180	M744	S341	M745
B392	B103	M746	S50	M747	S45	M748	S147	M749	S446	M750	S130	M751	S284	M752	S293	M753	S280	M754	S421	B390	S343	M755	B79
S440	B133	B60	M756	S54	M757	S245	M758	I608	M759	I615	B49	S259	I592	S303	M760	B230	I685	S328	M761	S90	M762	S241	M763
M764	S278	M765	S37	M766	B320	I696	S307	B25	S43	M767	S326	M768	S92	M769	S224	I680	S453	M770	S174	M771	S149	M772	M773
B168	I627	S176	M774	S69	M775	B362	M776	S186	M777	B190	I703	I590	B17	I586	I501	S222	I565	S402	M778	S410	M779	S182	M780
M781	S324	M782	S228	I498	S389	M783	S395	I563	S291	B135	B295	M784	S86	M785	S289	M786	S203	M787	S322	I656	I658	M788	S47
S264	M789	S404	M790	S301	B38	S470	M791	S466	B269	I562	M792	S132	B100	B299	M793	I676	I577	B99	B235	S382	M794	S406	I682
M795	S436	M796	S393	B267	I521				M797	I603	S457	I622	S376	B239				M798	B380	M799	B101	M800	I645
S432	M801	I497	M802	I530	I601				M803	S272	M804	S434	M805	M806				M807	M808	S138	I496	S414	B383
M809	S316	B179	S353	I513	M810				I570	M811	S30	I666	S261	M812				M813	S461	M814	I573	M815	B463
S48	M816	S305	M817	S276	M818				M819	I581	M820	S438	M821	I605				S16	M822	S159	M823	S178	I693
M824	I517	I667	S195	M825	S263				M826	B338	S397	M827	S337	M828				M829	S124	M830	S145	M831	S39
B400	I538	S197	I662	S213	M832				M833	S172	M834	S126	I526	B71				S80	M835	S270	M836	S73	M837
M838	S117	M839	B97	M840	S215				M841	M842	S449	M843	S417	M844				M845	S136	M846	I537	I523	S391
M847	B250	S345	M848	I525	M849				B32	S188	M850	S65	M851	B119				I594	I568	B163	S166	S232	M852
M853	B122	M854	B387	I515	S431				S56	M855	S161	B411	S41	M856				I683	I532	M857	S249	M858	M859
S312	B61	S355	M860	I712	M861				M862	I654	M863	S459	M864	S359				M865	M866	B419	B403	S58	M867
M868	S385	M869	I512	M870	S167				S24	M871	S33	I600	S128	M872				M873	S82	I520	S268	M874	S311
S88	M875	S297	M876	S451	M877				M878	S364	M879	S442	M880	B423				M881	M882	S20	M883	S199	M884
M885	I704	M886	S157	M887	S399				M888	M889	S234	M890	S349	M891				M892	S412	M893	S77	B91	I699
S120	M894	S226	M895	S444	B87				I602	S332	M896	B28	B177	M897				S216	M898	B427	M899	S318	M900
M901	S425	M902	S113	B300	I672				S408	M903	S366	M904	S314	M905				M906	I648	M907	I584	B34	S455
S472	M908	S368	M909	I719	B471				M910	S476	M911	S370	M912	M913				S464	M914	S374	M915	S474	M916

Section 4

Flag patterns


4.1 Protocols

4.1.1 Flag patterns, constant temperature


1. 85 °C for 3 minutes, no fluorescence measurement.
2. 71 °C to 55 °C, in steps of -2.0 °C every 4 minutes, taking a measurement at the end of each hold (36 minutes in total).
3. 53 °C to 47.2 °C, in steps of -0.2 °C every 12 minutes, taking a measurement at the end of each hold (6 hours in total).
4. A 47 °C hold, taking a measurement every 12 minutes (51 hours in total).
5. 47 °C to 40.5 °C, in steps of -0.5 °C every 13 minutes, taking a measurement at the end of each hold (3 hours 2 minutes in total).

4.1.2 Flag patterns, ramp

1. 71 °C for 4 minutes, no fluorescence measurement.
2. 71 °C to 55 °C, in steps of -2.0 °C every 4 minutes, taking a measurement at the end of each hold (36 minutes in total).
3. 53 °C to 48.2 °C, in steps of -0.2 °C every 12 minutes, taking a measurement at the end of each hold (5 hours in total).
4. 48 °C to 46.1 °C, in steps of -0.1 °C every 5 hours, taking a measurement every 30 minutes (100 hours in total).
5. 46 °C to 39.5 °C, in steps of -0.5 °C every 13 minutes, taking a measurement at the end of each hold (3 hours 2 minutes in total).




4.2 Nucleation model summaries





4.3 Individual results

4.3.1 H flag 1




4.3.2 H flag 2







4.3.3 H flag 3




4.3.4 H flag 4




4.3.5 H flag 5




4.3.6 H flag 6







4.3.7 H flag 7




4.3.8 H flag 8




4.3.9 H flag 9




4.3.10 H flag 10







4.3.11 H flag 11




4.3.12 H flag 12




4.3.13 A flag 1







4.3.14 A flag 2




4.3.15 A flag 3







4.3.16 A flag 4




4.3.17 A flag 5




4.3.18 A flag 6




4.3.19 A flag 7



4.3.20 A flag 8




4.3.21 A flag 9







4.3.22 A flag 10




4.3.23 A flag 11




4.3.24 A flag 12







4.3.25 M flag 1




4.3.26 M flag 2




4.3.27 M flag 3




4.3.28 M flag 4




4.3.29 M flag 5







4.3.30 M flag 6




4.3.31 M flag 7




4.3.32 M flag 8




4.3.33 M flag 9







4.3.34 M flag 10




4.3.35 M flag 11




4.3.36 M flag 12




4.3.37 M flag 13







4.3.38 Uniform concentrations (50 nM)




4.4 Plate-level fluorescence data used in Section 4.3

Kinetic nucleation control experiments, constant temperature, part 1






Kinetic nucleation control experiments, temperature ramp, part 1



Kinetic nucleation control experiments, temperature ramp, part 2




Section 5


Pattern recognition

5.1 Protocol

1. 71 °C for 4 minutes, no fluorescence measurement.
2. 71 °C to 55 °C, in steps of -2.0 °C every 4 minutes, taking a measurement at the end of each hold (36 minutes in total).
3. 53 °C to 48.2 °C, in steps of -0.2 °C every 12 minutes, taking a measurement at the end of each hold (5 hours in total).
4. 48 °C to 44.9 °C, in steps of -0.1 °C every 5 hours, taking a measurement every 30 minutes (160 hours in total).
5. 45 °C to 39.5 °C, in steps of -0.5 °C every 13 minutes, taking a measurement at the end of each hold (2 hours 36 minutes in total).



5.2 Nucleation model summaries



5.3 AFM shape counts summary

Counts of shapes were done manually by each of the four co-authors. Images were blinded by having a script assign random numbers to each image, and using PNG image files (with no AFM metadata) named using these numbers. CGE had previously seen all images associated with their corresponding samples while performing the AFM imaging.

Samples in the tables below are referred to as ‘H’, ‘A’, or ‘M’ for samples with four fluorophores on one shape, and ‘3’ for samples where each shape had one fluorophore (resulting in three total fluorophores). Each imaged sample (with a different fluorophore configuration) of each pattern had three $5 \mu\text{m} \times 5 \mu\text{m}$ images, with exception of the H, A, and M samples for Magnolia, which had two. To examine the potential effect of fluorophore and quencher tile modifications on shapes seen by AFM, several patterns had multiple samples imaged, while the ‘3’

sample was imaged for every pattern. Owing to the significantly larger numbers of shapes seen in the SHAM mix, only 1/4 of each image was used for counting. Numbers in the tables below are averaged across the counts from each co-author, and normalized to the number of shapes per μm^2 .

Shapes were classified into categories 1 and 2, where 1 were “almost complete” shapes, possibly missing small portions, while 2 were “clearly recognizable” shapes, which were distinctly not complete, but still clearly distinguishable. Detailed criteria distinguishing these categories were not developed: instead, each co-author interpreted the categories independently.

5.3.1 Counts averaged across co-authors and images, normalized to shapes per $25 \mu\text{m}^2$:


Pattern	Sample	H-1	H-2	A-1	A-2	M-1	M-2	total #	% target	%H	%A	%M
Hodgkin	3	4.17	41.00	0.00	0.00	0.00	0.92	46.08	98	98	0	2
Avogadro	3	0.00	0.00	8.25	21.25	0.00	0.75	30.25	98	0	98	2
Mitscherlich	3	0.00	0.17	0.00	0.08	14.00	36.92	51.17	100	0	0	100
Hopfield	3	13.92	30.08	0.00	0.00	0.00	0.33	44.33	99	99	0	1
Abbott	3	0.08	0.42	18.17	20.33	0.00	0.08	39.08	99	1	99	0
Moser	3	0.00	0.00	0.00	0.00	4.33	32.25	36.58	100	0	0	100
Horse	H	18.12	37.62	0.00	0.00	0.00	0.00	55.75	100	100	0	0
	A	22.08	55.50	0.00	0.00	0.00	0.00	77.58	100	100	0	0
	M	5.67	33.50	0.00	0.00	0.00	0.00	39.17	100	100	0	0
Anchovy	3	7.08	31.08	0.08	0.08	0.00	0.25	38.58	99	99	0	1
Mockingbird	H	0.83	4.75	0.00	0.00	1.25	4.92	11.75	52	48	0	52
	A	0.08	3.17	6.75	9.42	3.50	6.42	29.33	34	11	55	34
	M	0.00	2.25	0.00	1.17	1.58	10.58	15.58	78	14	7	78
	3	1.00	3.58	0.75	2.58	1.50	8.25	17.67	55	26	19	55
Hazelnuts	3	7.67	41.83	0.42	2.08	0.00	0.17	52.17	95	95	5	0
Apples	3	0.00	0.00	21.92	23.67	0.00	0.00	45.58	100	0	100	0
Magnolia	H	0.00	5.75	2.50	5.50	14.75	29.88	58.38	76	10	14	76
	A	0.00	3.00	9.25	10.38	23.25	45.50	91.38	75	3	21	75
	M	0.00	2.00	2.50	2.25	17.25	31.38	55.38	88	4	9	88
	3	0.00	0.25	0.33	0.42	3.25	17.25	21.50	95	1	3	95
Harom	3	5.33	47.83	0.08	1.25	0.08	1.17	55.75	95	95	2	2
Aon	H	0.00	1.67	3.17	8.42	1.25	4.17	18.67	62	9	62	29
	A	0.00	0.75	26.58	17.92	4.00	13.67	62.92	71	1	71	28
	M	0.25	2.75	16.75	23.12	4.00	20.12	67.00	60	4	60	36
	3	0.00	0.08	8.58	20.92	0.67	8.33	38.58	76	0	76	23
Mbili	H	0.42	8.75	1.83	4.17	22.75	29.83	67.75	78	14	9	78
	A	0.00	1.08	7.58	8.17	16.75	25.67	59.25	72	2	27	72
	M	0.00	0.83	0.42	0.75	10.67	17.58	30.25	93	3	4	93
	3	0.00	0.25	0.00	0.00	4.83	16.83	21.92	99	1	0	99
H	H	23.67	76.17	0.33	0.75	0.08	1.42	102.42	97	97	1	1
	A	4.33	36.92	2.00	3.17	0.00	1.17	47.58	87	87	11	2
	M	6.00	41.25	0.08	0.17	1.33	6.00	54.83	86	86	0	13
	3	0.67	26.17	0.00	0.25	0.17	1.67	28.92	93	93	1	6
A	3	0.00	1.42	19.17	17.83	0.33	4.00	42.75	87	3	87	10
M	3	0.00	0.33	1.50	4.58	14.50	19.67	40.58	84	1	15	84
NoisyApple1	3	0.08	0.08	0.08	0.42	0.08	0.08	0.83	60	20	60	20
NoisyApple2	3	0.00	0.00	0.00	0.67	0.00	0.42	1.08	62	0	62	38
NoisyApple3	3	0.00	0.00	2.67	10.42	0.00	0.17	13.25	99	0	99	1
NoisyHorse1	3	7.50	30.50	0.08	0.00	0.00	0.00	38.08	100	100	0	0
NoisyHorse2	3	0.00	0.50	0.00	0.00	0.00	0.00	0.50	100	100	0	0
NoisyHorse3	3	0.00	3.75	0.00	0.00	0.00	0.00	3.75	100	100	0	0
ObscMagn1	3	0.00	0.00	0.00	0.00	0.00	0.25	0.25	100	0	0	100
ObscMagn2	3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0	0	0	0
ObscMagn3	3	0.00	0.00	0.58	1.67	2.33	15.42	20.00	89	0	11	89
ObscHazel1	3	0.00	1.00	0.00	0.00	0.00	0.08	1.08	92	92	0	8
ObscHazel2	3	0.42	9.33	0.00	0.33	0.42	0.92	11.42	85	85	3	12
ObscHazel3	3	0.00	0.58	0.00	0.00	0.00	0.08	0.67	88	88	0	12
VarHarom1	3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0	0	0	0
VarHarom2	3	0.00	0.58	0.00	0.00	0.00	0.00	0.58	100	100	0	0
VarHarom3	3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0	0	0	0
VarHarom4	3	0.08	2.50	0.00	0.00	0.00	0.08	2.67	97	97	0	3
VarHarom5	3	0.00	1.08	0.08	0.58	0.00	0.17	1.92	57	57	35	9
VarHarom6	3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0	0	0	0
SHAM	H	53.75	164.08	48.25	47.00	44.92	50.00	408.00	100	53	23	23
	A	7.83	67.17	123.14	53.75	43.64	38.44	333.97	100	22	53	25
	3	18.58	92.42	48.08	47.08	64.75	35.33	306.25	100	36	31	33

5.3.2 Counts averaged across co-authors, images and samples, normalized to shapes per 25 μm^2 :


Pattern	H-1	H-2	A-1	A-2	M-1	M-2	total #	% target	%H	%A	%M
Hodgkin	4.17	41.00	0.00	0.00	0.00	0.92	46.08	98	98	0	2
Avogadro	0.00	0.00	8.25	21.25	0.00	0.75	30.25	98	0	98	2
Mitscherlich	0.00	0.17	0.00	0.08	14.00	36.92	51.17	100	0	0	100
Hopfield	13.92	30.08	0.00	0.00	0.00	0.33	44.33	99	99	0	1
Abbott	0.08	0.42	18.17	20.33	0.00	0.08	39.08	99	1	99	0
Moser	0.00	0.00	0.00	0.00	4.33	32.25	36.58	100	0	0	100
Horse	12.80	39.59	0.02	0.02	0.00	0.07	52.50	100	100	0	0
Anchovy	0.00	0.33	20.33	23.17	0.00	0.08	43.92	99	1	99	0
Mockingbird	0.48	3.44	1.88	3.29	1.96	7.54	18.58	51	21	28	51
Hazelnuts	7.67	41.83	0.42	2.08	0.00	0.17	52.17	95	95	5	0
Apples	0.00	0.00	21.92	23.67	0.00	0.00	45.58	100	0	100	0
Magnolia	0.00	2.47	3.28	4.17	13.36	29.47	52.75	81	5	14	81
Harom	5.33	47.83	0.08	1.25	0.08	1.17	55.75	95	95	2	2
Aon	0.05	1.18	13.50	17.09	2.34	10.80	44.95	68	3	68	29
Mbili	0.10	2.73	2.46	3.27	13.75	22.48	44.79	81	6	13	81
H	8.67	45.12	0.60	1.08	0.40	2.56	58.44	92	92	3	5
A	0.00	1.42	19.17	17.83	0.33	4.00	42.75	87	3	87	10
M	0.00	0.33	1.50	4.58	14.50	19.67	40.58	84	1	15	84
NoisyApple1	0.08	0.08	0.08	0.42	0.08	0.08	0.83	60	20	60	20
NoisyApple2	0.00	0.00	0.00	0.67	0.00	0.42	1.08	62	0	62	38
NoisyApple3	0.00	0.00	2.67	10.42	0.00	0.17	13.25	99	0	99	1
NoisyHorse1	7.50	30.50	0.08	0.00	0.00	0.00	38.08	100	100	0	0
NoisyHorse2	0.00	0.50	0.00	0.00	0.00	0.00	0.50	100	100	0	0
NoisyHorse3	0.00	3.75	0.00	0.00	0.00	0.00	3.75	100	100	0	0
ObscMagn1	0.00	0.00	0.00	0.00	0.00	0.25	0.25	100	0	0	100
ObscMagn2	0.00	0.00	0.00	0.00	0.00	0.00	0.00	nan	nan	nan	nan
ObscMagn3	0.00	0.00	0.58	1.67	2.33	15.42	20.00	89	0	11	89
ObscHazeln1	0.00	1.00	0.00	0.00	0.00	0.08	1.08	92	92	0	8
ObscHazeln2	0.42	9.33	0.00	0.33	0.42	0.92	11.42	85	85	3	12
ObscHazeln3	0.00	0.58	0.00	0.00	0.00	0.08	0.67	88	88	0	12
VarHarom1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	nan	nan	nan	nan
VarHarom2	0.00	0.58	0.00	0.00	0.00	0.00	0.58	100	100	0	0
VarHarom3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	nan	nan	nan	nan
VarHarom4	0.08	2.50	0.00	0.00	0.00	0.08	2.67	97	97	0	3
VarHarom5	0.00	1.08	0.08	0.58	0.00	0.17	1.92	57	57	35	9
VarHarom6	0.00	0.00	0.00	0.00	0.00	0.00	0.00	nan	nan	nan	nan
SHAM	26.72	107.89	73.16	49.28	51.10	41.26	349.41	100	39	35	26

5.4 Individual fluorescence and AFM results


5.4.1 Hodgkin




5.4.2 Avogadro




5.4.3 Mitscherlich




5.4.4 Hopfield




5.4.5 Abbott




5.4.6 Moser




5.4.7 Horse







5.4.8 Anchovy




5.4.9 Mockingbird







5.4.10 Hazelnuts




5.4.11 Apples




5.4.12 Magnolia







5.4.13 Harom





5.4.14 Aon







5.4.15 Mbili






5.4.16 LetterH







5.4.17 LetterA




5.4.18 LetterM




5.4.19 NoisyApple1




5.4.20 NoisyApple2




5.4.21 NoisyApple3




5.4.22 NoisyHorse1




5.4.23 NoisyHorse2




5.4.24 NoisyHorse3




5.4.25 ObscMagn1




5.4.26 ObscMagn2




5.4.27 ObscMagn3




5.4.28 ObscHazeln1




5.4.29 ObscHazeln2




5.4.30 ObscHazeln3




5.4.31 VarHarom1




5.4.32 VarHarom2




5.4.33 VarHarom3




5.4.34 VarHarom4



5.4.35 VarHarom5



5.4.36 VarHarom6



5.4.37 Uniform concentrations (60 nM)

